

Dark Passage: Mental Health Consequences of Parental Death*

Petri Böckerman, University of Jyväskylä, Labour Institute for Economic Research, and IZA Institute of Labor Economics. E-mail: petri.bockerman@labour.fi

Mika Haapanen, University of Jyväskylä, School of Business and Economics. E-mail: mika.p.haapanen@jyu.fi

Christopher Jepsen, University College Dublin, School of Economics and Geary Institute, IZA Institute of Labor Economics, and CES-Ifo. E-mail: christopher.jepsen@ucd.ie

9th September 2021

Abstract:

This paper studies the causal effect of parental death on children's mental health. Combining several nationwide register-based data for Finnish citizens born between 1971 and 1986, we use an event study methodology to analyze hospitalization for mental health-related reasons by the age of 30. We find no clear evidence of increased hospitalization following the death of a parent of a different gender, but there are significant effects for boys losing their fathers and girls losing their mothers. Depression is the most common cause of hospitalization in the first three years following paternal death, whereas stress and, to a lesser extent, self-harm are the most common causes five to ten years after paternal death. Using family fixed effects models, the largest negative effects on mental health prevail for those boys who lose their father.

Keywords: parental death, mental health, hospitalization, depression, sickness absence

JEL classification codes: I10, I12, J12, J13

* This study is part of a project supported by Yrjö Jahnsson Foundation (no. 6621), and OP Group Research Foundation. We thank Christian Hakulinen, Yannick Dupraz, and seminar participants at the Annual Conference of the Royal Economic Society (2021), the Annual Conference of the European Society for Population Economics (2021), Irish Economic Association Annual Conference (2021), University College Dublin, Essen Health Conference 2021, Social Insurance Institution of Finland (Kela), the Finnish Institute for Health and Welfare (THL), and Labour Institute for Economic Research, for their useful comments.

Introduction

Children face many challenges on their path to adulthood. Probably the most difficult situation a child faces is the death of a parent, which is often the first time that a child deals with death, sorrow, and ordeal. In this critical point in life, children are forced to encounter and overcome a loss of parental guidance and social support, a likely reduction in family income and economic resources, and many other shortfalls that other children do not have to face. The traumatic event likely leaves scars across a host of outcomes such as future educational outcomes, labor-market outcomes, family formation, and health.

Mental health is an increasingly important determinant of overall health in all developed countries (Layard, 2013), with depression being the largest contributor to the disease burden attributable to non-fatal health outcomes (Whiteford et al., 2013). Mental disorders at young ages have an enduring impact on individuals' wellbeing and economic prosperity. Therefore, early mental health-related problems often accumulate into negative health and non-health consequences in adulthood. For example, mental health problems may lead to poor physical health (Sareen et al., 2006) as well as work-related losses, such as lower work performance and increased absenteeism (Bubonya, Cobb-Clark, and Wooden, 2017). Finally, individuals with severe mental illness die on average 10-20 years earlier than the population in general (Liu et al., 2017).

Despite the increasing importance of mental health, economic research has only rarely analyzed the effect of parental death on the child's mental health. To gain a better understanding of the drivers of mental health differences in the population, the preferred outcome of our analysis is hospitalization for mental health-related reasons. We also examine other closely related outcomes such as the use of mental health-related medications and sickness absence. Our analysis is based on nationwide register-based data from Finland for individuals born between 1971 and 1986, in order to measure hospitalization through the age

of 30. Population-based data are free from health-related selection and non-participation biases.

For identification, our preferred specification is an event study framework akin to the model used by Kleven, Landais, and Egholt Søgaard (2019) and Kristiansen (2021), in which we follow individuals before and after a parental death at ages 10–20. An extension to the event study model uses children who did not experience a parental death as a comparison group. To check the sensitivity, we estimate models based on deaths driven only by likely random causes and not correlated with socioeconomic status. To account for time-invariant characteristics of the parental or family background, we also present results using family fixed effects. Our supplementary analysis uses a cross-sectional model based on additional outcomes such as sickness absence, employment, and earnings in adulthood.

Strikingly, based on the preferred event study models, we find that losing a parent of one's own gender has a much larger negative effect on mental health, than losing a parent of the opposite gender. For males, the pre-death hospitalization rate doubles after the loss of the father. Although the effect diminishes in later years, it is still economically and statistically significant three years after the death. The overall effect of the mother's death on females fades noticeably faster. We find little if any evidence of adverse mental health outcomes associated with the mother's death for males or the father's death for females. This pattern of results is similar, whether or not we use children who did not experience a parental death as a control group. Looking at the causes of hospitalization, depression is the most pronounced cause in the short run, and stress in the long run among males, after a paternal death. Further, we find substantial increases in the use of mental health-related medications, sickness absence and employment/earnings losses for the affected children in adulthood.

Relationship to Previous Work

Originally, economists studied parental death because they considered it the most exogenous source of parental absence. In a meta-analysis, Amato (1993) suggests that parental divorce

had larger negative consequences than parental death. In contrast, Corak (2001) finds few differences between parental death and parental divorce in multiple labor-market outcomes in an analysis of changes in Canadian divorce laws.

Two early studies look simultaneously at the effects of parental death and parental divorce, using OLS models on rich, cross-sectional survey data. In regressions that also include the number of years living with each parent, Lang and Zargosky (2001) find no discernable effect of parental death on numerous economic outcomes in adulthood for National Longitudinal Survey of Youth data. Similarly, Fronstin, Greenberg, and Robins (2001) study both parental divorce and father's death (data on mother's death were not collected) on outcomes at age 33, using the British National Child Development Survey. Looking either at the effect of death overall or at the effect at different ages of the child, they find no systematic relationship between father's death or parental divorce on labor-market status or earnings by gender.

More recently, Steele, Sigle-Rushton, and Kravdal (2009) look at both marital dissolution (separation or divorce) and father's death using Norwegian administrative data. To account for potential selection into marital dissolution, they use a simultaneous equations model with separate models for marital dissolution and for children's education. In their preferred model accounting for selection, they find similar negative impacts for marital separation and father's death on the transition from lower to higher secondary education.

Chen, Chen, and Liu (2009) use detailed registry data from Taiwan to study the effect of accidental parental death on the likelihood of college enrollment at age 18. Their preferred model is a family fixed effects model that compares siblings who were 18 or more when the parent died, with younger siblings. They find that any maternal death is associated with a two-percentage-point reduction in college attendance, compared to a reduction of more than four percentage points for accidental maternal deaths. The impact of paternal death is small and insignificant. Because both siblings are probably near 18 in age, Adda, Björklund, and

Holmlund (2011) note that results from this technique likely do not generalize to the entire population of parental deaths. Kailaheimo-Lönnqvist and Erola (2020) also utilize family fixed effects models and find negative impacts of early parental death on children's university education, in Finland.

Also using registry data, Adda, Björklund, and Holmlund (2011) study the relationship between parental death, education, and earnings. They report negative effects of parental death, with more pronounced effects for mothers' death in OLS regressions, either for all deaths or as a subset of plausibly exogenous causes of death. When they account for selection using a technique similar to selection on observables versus unobservables (Altonji, Elder, and Taber, 2005), most of these effects become insignificant.

Gimenez et al. (2013) consider the impact of plausibly exogenous causes of parental death in Taiwan, finding that parental death has a negative effect on educational attainment, coupled with a positive effect on the likelihood of entering the labor force before age 20. Females experiencing a parental death are more likely to marry by age 20, whereas males experiencing a parental death are more likely to enlist in the military within four years of completing secondary school.

Investigating the relationship between parental death and children's occupational status using historical data (1850 to 1952) from the Netherlands, Rosenbaum-Feldbrügge (2019) finds, for both men and women, larger adverse effects due to maternal loss compared to paternal loss in OLS models. Similarly, Dupraz and Ferrara (2021) document a negative effect of paternal death for men whose fathers served in the Union Army during the U.S. Civil War.

Our paper is most closely related to the concurrent work by Kristiansen (2021),¹ who examines the effect of parental health shocks, including deaths, on the likelihood of two

¹ Less closely related, but still relevant, is the work by Persson and Rossin-Slater (2018), who report the adverse mental health consequences for children whose mothers experienced a death in the family while pregnant.

mental health-related outcomes, therapy and anti-depressant medication. She studies health shocks for children aged 14 to 18 at the time of the shock, using Danish registry data. Based on event study framework, she finds that parental death is associated with short-run increases in both outcomes, with therapy more likely among higher-income families and anti-depressants among lower-income families. In the long run, anti-depressant use is correlated with lower education, whereas no such association is evident for therapy.

Rather than analyzing the direct impact of parental death on child outcomes, Kalil et al. (2016) and Gould, Simhon, and Weinberg (2019) investigate how parental death changes the relationship between parental schooling and children's education outcomes in Norway and Israel, respectively. They find that the educational level of the parent who dies becomes less important for children's education outcomes after the parent's death.

A substantial body of literature on the effects of parental death exists outside economics, using mainly cross-sectional estimation strategies that do not permit causal claims about the estimated relationships. However, reviewing that literature is beyond the scope of the current work. Instead, we highlight few examples and encourage interested readers to see them and the citations within. In epidemiology, Appel et al. (2013) use a hazard model to estimate the association between parental death and the risk of hospitalization for affective disorders such as bipolar disorder and schizophrenia. They use registry data from Denmark through 2009. A similar study in psychology (Berg, Rostila, and Hjern, 2016) looks at parental loss and depression in Sweden, while McKay et al. (2021) provide an up-to-date meta-analysis of studies in public health.

We contribute to the literature by analyzing the causal relationship between parental death and mental health-related hospitalization. We differ from previous work, particularly from Kristiansen's (2021) analysis of parental death in Denmark that studies therapy and anti-depressant medication as outcomes, in five ways. First, we focus on mental health-related hospitalization and estimate separate effects by parent's gender. Second, to identify

the explicitly exogenous causes of death, we apply the approach of Espinosa and Evans (2008) and Gimenez et al. (2013), which permits us to estimate the causal effects using deaths not correlated with parental socioeconomic status (i.e., income and education). Third, we present results using family fixed effects accounting for time-invariant characteristics of the parental or family background. Fourth, we provide a second event study analysis, using a difference-in-differences framework, by including a comparison group of individuals without a parental death. Fifth, we test more formally the underlying assumption of parallel trends, using the techniques of Borusyak et al. (2021). The event study analysis used here and in Kristiansen (2021) explicitly focuses on how parental death affects mental health over time, in contrast to previous works where the outcome is typically defined only at a specific point in time, such as a college entrance exam (Chen, Chen, and Liu, 2009; Gould, Simhon, and Weinberg, 2019).

Administrative datasets

Our empirical analysis is based on nationwide administrative data sources. We start by describing health registers and proceed to characterize the census data. Finally, we describe the outcome variables and provide key descriptive information on parental death.

Health Registers

We use data from the comprehensive death certificates compiled by Statistics Finland to identify the cause and date of death over the period 1970–2016. All diagnoses for the causes of death pass a routine validation conducted by Statistics Finland, and unclear cases are judged by a panel (Lahti and Penttilä, 2001).²

The main source of mental health data is the Discharge Register from the Finnish Institute for Health and Welfare, which identifies all inpatient discharges in specialized

² The statistics on causes of death include all deaths in Finland or abroad of persons permanently resident in Finland at the time of their death.

public health care for the Finnish population over 1970–2016. Outpatient visits to specialized mental health care facilities are recorded over a shorter period of 1998–2016. In typical cases, several diagnostic procedures have contributed to the diagnosis, including an additional structured clinical interview in some cases. Diagnoses for mental-health disorders are usually established by several treating doctors also.³ Finland’s national health insurance system covers all citizens, with almost all hospitalizations in the public sector, with a very small private Finnish health care system, providing outpatient care principally.

We focus on hospitalization as the main outcome of our analysis for four reasons. First, the treatment costs of mental health-related hospitalizations are considerable in the universal health care system. Second, serious mental illnesses cause considerable indirect economic costs, in terms of weak long-run labor market attachment and lost earnings over the course of life (Hakulinen et al., 2019). Mental disorders are also the leading cause of disability pensions in Finland. Third, the overall reliability of hospitalization data for empirical research is well established and the measurement error very small (Sund, 2012). Fourth, data are available for an extensive period from 1970 onwards, facilitating the use of event study framework to analyze the dynamic effects.

The additional health outcomes (e.g., sickness absenteeism and the usage of mental health-related medications) used in cross-sectional models are described in Appendix D.

Census Data

These health registers are linked to the census data on the population of Finland, available from Statistics Finland. The census files are available at five-year intervals from 1970 to 1985 and annually from 1987 to 2016 and provide comprehensive information on the parents and their children, including data on family composition, education, earnings, occupation, and the region of residence.

³ Because hospitalization involves multiple steps, any counseling or other services provided to children after a parental death is unlikely to produce a ‘mechanical’ increase in hospitalization.

Given that Finland has a current population of approximately 5.5 million, we use data for an extended time period, for both parents and children, to improve the precision of the estimates. Specifically, we follow Finnish individuals born between 1971 and 1986, by which we have data on approximately one million individuals who have reached at least the age of 30 (in 2016). We exclude children born outside Finland and those who have no data for either parent.

Outcome Variables

With these linked data,⁴ we analyze several outcome variables. Event study framework requires observations over a long time to identify the dynamic adjustment to parental death. Therefore, our main outcome of analysis using an event study approach describes whether an individual had at least one (in-patient) hospitalization spell annually due to mental health-related disorders (ICD-10: F, ICD-8 and ICD-9: 290–319). To obtain a more fine-grained picture, we also examine selected conditions related to i) depression, ii) stress, and iii) substance-use disorder. Further, we study effects on hospitalizations due to self-harm (including suicide attempts). In the hospitalization records, self-harm attempts (ICD-10: X60–X84, ICD-8 and ICD-9: E950–E959) are recorded as independent, external causes of hospitalization (i.e., not all self-harming attempts are coded as mental health disorders). Of self-harming attempts that lead to hospitalization, mental health disorder is recorded as the principal cause of hospitalization in approximately 10% of the cases.

In the cross-sectional analysis, we utilize several other outcomes such as a broader measure of mental health-related deaths, sickness absence, employment, and earnings in adulthood, for which we do not have observations over the long period that would allow us to examine the dynamic effects using event study framework. Following Alexander and Schnell (2019), we analyze a broader measure of mental health-related deaths as additional outcome.

⁴ As the linkage between data sources is done based on a unique person identifier akin to the Social Security number in the U.S., the data – as in other Nordic countries – are of very high quality.

The measure includes not only officially recorded suicides in death certificates, but also injuries of undetermined intent (i.e., fatal injuries unascertainable as to whether they were accidental or purposely inflicted), and accidental deaths involving poisonings, drownings, and deaths involving firearms and trains. We prefer this more comprehensive measure of mental health-related deaths to suicides because all deaths caused by mental health-related disorders are not necessarily classified officially as suicides in the register data.

In addition to the hospital spells in inpatient care, we also observe whether individuals had day visits to special care units related to mental health disorders at ages 27–30. We further measure the number of sickness absence days (ages 26–30), whether they have sickness absence spells due to mental health reasons (age 30) and whether they use mental health-related medications (ages 29–30). To analyze the potential mechanisms at play, we analyze the effects on children’s employment and earnings (at ages 26–30), based on comprehensive information from the Finnish tax authorities, and years of schooling (by age 30), based on register-based information on completed degrees available from Statistics Finland since 1970.

Descriptive Information on Parental Death

Parental death during childhood and early adulthood is relatively rare in developed countries. In our data approximately 15 percent of individuals experience a parental death before they turn 31 (Table 1). We observe two additional patterns. First, parental death experiences in childhood and early adulthood are skewed towards the death of the father. Less than five percent of individuals before they turn 31 experience the death of their mother, compared to nearly 12 percent for the death of the father. This pattern presumably leads to less precise estimates for the impact of maternal deaths. Second, for either parent, the likelihood of death increases substantially with the individual’s age, from under one percent for a parental death before age 10 to 4.64 percent when the individual is between 26 and 30 years old.

Table 1: Age of Individual upon Parent's Death

| Age when Parent Died | Death of Mother | | Death of Father | | Death of Parent | |
|----------------------|-----------------|---------|-----------------|---------|-----------------|---------|
| | Freq. | Percent | Freq. | Percent | Freq. | Percent |
| Age 0–2 | 714 | 0.07 | 2,819 | 0.30 | 3,513 | 0.36 |
| 3–6 | 1,595 | 0.17 | 5,802 | 0.61 | 7,332 | 0.76 |
| 7–10 | 2,573 | 0.27 | 8,093 | 0.85 | 10,496 | 1.09 |
| 11–15 | 4,853 | 0.51 | 14,109 | 1.49 | 18,500 | 1.92 |
| 16–20 | 7,378 | 0.77 | 19,728 | 2.08 | 26,059 | 2.70 |
| 21–25 | 10,434 | 1.09 | 26,633 | 2.81 | 34,898 | 3.62 |
| 26–30 | 14,293 | 1.49 | 35,096 | 3.70 | 45,212 | 4.69 |
| No death by age 30 | 917,778 | 95.64 | 836,725 | 88.17 | 818,923 | 84.87 |
| Total | 959,618 | 100.0 | 949,005 | 100.0 | 964,933 | 100.0 |

Notes: Number of observations is smaller for fathers because it is more common that the link between parent and child is missing for father (1.7%) than mother (0.6%).

Methods

Event Study Specifications

To allow the relationship between parental death and children's mental health outcomes to vary with the time since death, our main results are based on an event study specification, analogous to the model estimated by Kleven et al. (2019) for the effect of children on gender inequality. This approach allows the analysis of dynamics and adaptation to the shock with an emphasis on the length of the effects detected on mental health.

Equation (1) depicts the event study specification:

$$Y_{ist} = \sum_{j \neq -1} \alpha_j \cdot I[t = j] + \sum_k \beta_k \cdot I[age_{is} = k] + \sum_y \tau_y \cdot I[s = y] + \varepsilon_{ist} \quad (1)$$

The preferred measure of Y_{ist} is a dummy variable that equals to one for person i being hospitalized for a mental health-related condition at age s in time t (year relative to parental death).⁵ The right-hand side of the equation captures the events through a series of indicators (or dummy variables) for each of the three different aspects of the event. The first set of coefficients (α_j) captures the effect of the parental death at time $t = j$. Specifically, we include fixed effects for each year from eight years before the parental death until ten years

⁵ See Figure A1 in the Supplementary Appendix for the development of hospitalization rates before and after the parental death in the treatment groups.

after the parental death. The second set of coefficients (β_k) controls for the effects of the age of the child, thereby capturing age-specific vulnerability to mental health problems leading to hospitalization. Finally, the third set of coefficients (τ_y) accounts for calendar year effects. To capture the potential heterogeneity in the effects, we estimate separate event study models for male and female children, as well as for maternal and paternal death.

This event study model is estimated only for individuals who had experienced a parental death. In fact, to allow for effects up to eight years prior to and ten years after the parental death, the sample for the event study models is limited to individuals who were 10 to 20 when a parent died.⁶ To infer a causal effect from an event study model, children who are treated at different ages should have similar trends in pre-parental-death hospitalization. Following Borusyak et al. (2021), we examine this key identification assumption by conducting formal statistical tests of the pre-parental-death trends by using data from the pre-treatment periods. The results reported in Appendix B do not show evidence of significant pre-trends (based on individual t-tests or a joint significance F-test; see Table B1).

Our second model is a difference-in-differences event study specification, again following the estimation technique of Kleven et al. (2019). Specifically, we create a control group of individuals who did not experience a parental death by assigning pseudo death years for their father (and mother) in such a way that it follows the same (conditional) discrete distribution (for the age of death) as deaths in the treatment group.⁷ Separate event study regressions, as described in equation (1), are estimated for the treatment and control groups, in addition to separate models by parental death and gender of the child. Here, the assumptions required for causality are that the trends in pre-parental-death hospitalization are similar between children who experienced a parental death and those who did not. This

⁶ In Kristiansen (2021), the age range is 14 to 18, and the time period is from four years prior to and five years after the parental health shock.

⁷ Note that Kleven et al. (2019) use log-normal distribution for the births; but we cannot use it here, because the deaths are clearly not following normal or log distribution (when the child is 20 or under).

assumption is supported by Tables B1 and B2 that do not show evidence for significant pre-trends in the treatment or control group.

We also examine the robustness of our baseline results using the imputation estimator proposed by Borusyak et al. (2021).⁸ Contrary to fixed effects regression with lags and leads of treatment, the coefficients from the imputation estimator are robust in the presence of heterogeneous treatment effects (see Appendix B).⁹

The main concern regarding the causal interpretation of our baseline estimates is that some causes of death are arguably more exogenous than others. For instance, the medical literature has shown that cardiovascular diseases are strongly correlated with socioeconomic status (SES), a lower SES being significantly associated with higher cardiovascular risk (Tousoulis et al., 2020). We address this endogeneity concern in two ways. First, to identify the explicitly exogenous causes of death, we use the approach introduced in Espinosa and Evans (2008) and adopted later in Gimenez et al. (2013). The basic idea of this approach is to classify the causes of death into two groups: i) deaths strongly correlated with measures of parental socioeconomic status (i.e., informative causes of death) and ii) deaths driven by likely random causes and not correlated with socioeconomic status (i.e., uninformative causes of death). This classification implies that uninformative causes of death are, by construction, unrelated to parental socioeconomic characteristics and provide a clean source of exogenous variation in parental death. Thus, to check the robustness of our baseline findings, we estimate event study models separately for uninformative and informative causes of death. The empirical implementation of this approach is described in Appendix C.

The second approach to address potential endogeneity of parental death is a family fixed effects model as in Chen, Chen, and Liu (2009). This approach is based on the stylized fact that according to the literature, parental death tends to be more harmful the younger the

⁸ See von Bismarck-Osten et al. (2020) for an application of the method.

⁹ Besides the method proposed by Borusyak et al. (2021) there are also other methods to address the potentially heterogeneous treatment effects in the event study framework. However, the results in Freyaldenhoven et al. (2021, p. 32) suggest that these methods tend to produce largely similar time profiles for the effects.

child is (e.g., Kailaheimo-Lönnqvist and Erola, 2020), as parental influence is likely to diminish over time. By comparing the outcomes of same family children – and in some cases, the same gender – who lose their parent at a younger versus older age, we evaluate the effect of the parental death, eliminating time-invariant characteristics of the family. Therefore, between-sibling differences effectively differentiate sibling-invariants observed and unobserved confounding variables, including potential genetic and environmental influences. For example, parents may at least to some extent self-select into the increased likelihood of premature death by engaging in risky behaviors such as alcohol consumption and smoking due to genetic and environmental factors, which can also be correlated with mental health-related disorders.

Cross-Sectional Specifications

As some outcomes are available only in adulthood, we also present descriptive results based on a cross-sectional ordinary least squares (OLS) regression, as in Rosenbaum-Feldbrügge (2019):

$$Y_i = F_i\alpha_f + M_i\alpha_m + X_i\beta + \varepsilon_i \quad (2)$$

As before, the preferred outcome (Y_i) is a dummy variable that equals one for individuals ever hospitalized for a mental health-related condition at ages 26–30. Here, F_i and M_i represent a set of dummy variables for different ages of death (0–6, 7–10, 11–15, 16–20, 21–25, or 26–30) for the father (F) and mother (M), respectively. The comparison group is the set of individuals who did not experience a parental death by age 30. X_i is a set of control variables including mother’s and father’s education, income, occupation and mental health, native language, mother’s age at childbirth, number of siblings, as well as birth year and birth region fixed effects.¹⁰

¹⁰ Appendix Table D1 provides descriptive statistics at the individual level, separately by parental death. Strikingly, the rate of hospitalization for mental health-related disorders at ages 26–30 is 2.0 percent for individuals with no parental death by age 30, compared to rates of 3.3-3.7 percent for individuals with a parental death by age 30.

Controlling for parental education and income is important because socioeconomic status (SES) is correlated with longevity, i.e., children with lower SES parents are more likely to lose them early. A strong identifying assumption in this model is that the set of control variables (X_i) captures all the underlying differences between children with a parental death and children without a parental death. In a robustness test, we also report the main cross-sectional results for the uninformative causes of death, where the identifying assumptions are more likely to hold.

The major difference between the baseline event study model and cross-sectional model is that the event study model compares hospitalization rates (within-person) before and after parental death. In contrast, the cross-sectional model compares hospitalization rates at ages 26–30 between individuals who experience parental death at different ages. In other words, the event study model represents the timing of the short-term impact, whereas the cross-sectional model represents the cumulative, long-run impact. This difference in timing and the different underlying identification assumptions preclude any direct comparison of the results from the two estimation techniques.

For both the event studies and the cross-sectional regressions, all results are based on linear probability models, even though the preferred outcome is binary. These models facilitate the easier interpretation of the estimated coefficients and are also less sensitive to distributional assumptions.

Results


The Main Event Study Estimates

We start our analysis by presenting the estimates from our event study model based on equation (1). The results from this model are shown graphically in Figure 1, and the estimates are reported in Table A1. This model uses panel data for individuals who had a parental death when they were between 10 and 20 years, with yearly observations from eight years before

the death up to ten years after the death (i.e., up to 19 observations per person). The dependent variable is a dummy variable equal to one for individuals who are hospitalized with a mental health condition in the year. The reference period is the year prior to parental death when the average hospitalization rate was between 0.004 and 0.006. The figure plots the change in likelihood of hospitalization relative to the year before the parental death.

Figure 1 shows that males and females have a large increase of 0.004 in the likelihood of hospitalization for mental health conditions in the year the parent of the same gender dies, compared to the year prior to the parental death. For females, this effect declines quickly to 0.0028 and 0.0024 in the next two years. Thereafter, the coefficients are not statistically significant from zero at the ten-percent level for a two-sided test. For males, the coefficients for paternal death decline more slowly, with a statistically significant effect of 0.0030 three years after the death. Of the 11 coefficients for zero to ten years after the death, five are significant at the one-percent level, three at the five-percent level, two at the ten-percent level, and one is not statistically significant at the ten-percent level (all tests are two-sided). In other words, the effect of a paternal death for a male aged 10 to 20 is strongest in the short term, but sizable effects persist in the long term, too.

Figure 1: Event Study Results Using Hospitalization for Males and Females



Notes: The figures plot the coefficient estimates from event study regressions, together with 95% confidence intervals (standard errors clustered at the individual level). The outcome is the hospitalization rate. Panels on the left show estimates for father’s death and those on the right show them for mother’s death.

By comparison, the effect of a parent of a different gender dying is much smaller. For males, the effect of a maternal death is 0.0026 and marginally significant in the year of the death, followed by a statistically insignificant coefficient of 0.0020 in the year after the death. After that, the coefficients are generally close to zero and are never statistically significant at the ten-percent level. None of the coefficients for females experiencing paternal death are statistically significant at the ten-percent level, and only two of them are above 0.001 in magnitude.

Past work focusing on educational and labor-market outcomes has generally found that the largest effects are for maternal death (Rosenbaum-Feldbrügge, 2019; Chen, Chen, and Liu, 2009). Although we find short-term effects of maternal death for females, our largest (cumulative) effects on mental health are for males experiencing a paternal death. This result

is broadly consistent with the lower probability of marriage among males experiencing a paternal death found by Lang and Zagorsky (2001), as well as larger negative effects of paternal death on non-cognitive outcomes in Adda, Björklund, and Holmlund (2011).¹¹

The broader social science literature is also helpful for providing additional interpretation for our main finding that the death of a parent of a different gender has much smaller effects on hospitalization. First, a child of the same sex might be expected to take the lost parent's role in the family. Unexpected additional responsibilities at young ages could lead to a substantial amount of mental strain and stress (e.g., after the death of a mother the daughter may be forced to do more of the housework and take care of well-being of younger siblings in the family). Second, emotional attachment may be stronger if the parent and child have the same sex. For example, psychological literature has provided evidence that fathers tend to spend more shared time with their sons than with daughters (Raley and Bianchi, 2006), which strengthens the psychological ties between fathers and sons.

Gender preferences and role models may also play a role. Fathers may invest more economic and non-economic resources to boys than daughters. In contrast, mothers may prefer daughters over boys.¹² In addition, the parent of the same sex is arguably an important role model for the development of child's personality traits and the provider of social support for the child (e.g., Wiese and Freund 2001; Bokhorst et al., 2010). For example, the lack of social support may cause difficulties in school. Thus, the negative mental effects are arguably larger when the parent of the same gender dies.

¹¹ An earlier Finnish study supports the view that boys are more vulnerable to shocks at young ages in terms of mental health. Using the separations that took place during World War II when Finnish children were voluntarily evacuated unaccompanied by their parents to temporary foster care abroad Rääkkönen et al. (2011) show that the separated men showed a higher risk of any mental and substance use disorders than the non-separated men later in life. In contrast, separated and non-separated women did not differ from each other in the risk of mental disorders.

¹² The evidence from Finland regarding parental gender preferences shows that in the 1970s and early 1980s there was a parental boy preference in the families (Saarela and Finnäs, 2014). However, parental boy preference has been practically non-existent since the 1990s.

Robustness of the Main Event Study Estimates

Next, we evaluate the robustness of the baseline event study results to alternative specifications. In addition to the results for our baseline event study model (columns 1 and 4), Tables A2a–A2b report the results for a specification with the addition of individual fixed effects (columns 2 and 5), and a model with the additional control variables as in the cross-sectional model (columns 3 and 6). The tables show that all the results are quite consistent across these three models for males.¹³ Similarly, Figure B1 shows the similarity of results when using the imputation estimator proposed by Borusyak et al. (2021), a more flexible event study framework allowing for heterogeneous treatment effects.

Tables A3a–A3d present results by cause of hospitalization (see also Figures A2a, A2b, A3). In each table, the first column reports the result from Table A1 for all causes, and the remaining columns present the results for a specific cause of hospitalization: (2) depression, (3) stress, (4) substance abuse and (5) intentional self-harm (incl. suicide attempts).¹⁴ For males experiencing a paternal death, the coefficients are largest for depression in the short term and for stress in the long term. For males experiencing a maternal death, the results are inconclusive: the estimated post-parental-death coefficients are imprecise, and several pre-parental-death coefficients for stress are marginally significant.

For females, the most pronounced effects for maternal death are associated with intentional self-harm and depression. At the same time, there are marginally significant effects six to seven years before the death, suggesting some caution in attributing much emphasis to the post-death effects. For females experiencing a paternal death, the estimated coefficients are mostly small and statistically insignificant. As these hospitalization outcomes

¹³ A relatively minor exception to this pattern is that, for females, the coefficients for father's death are statistically significant in one of the three models. We adopt a conservative attitude in the interpretation of the results and conclude that there is no consistent relationship between father's death and daughter's mental health outcomes, a result in agreement with the insignificant results for mother's death and son's mental health outcomes.


¹⁴ This list is not exhaustive. For space considerations, we have not included results for schizophrenia or other rare mental health conditions. The prevalence of schizophrenia is also very low, making it difficult to identify statistically significant effects; further, it is largely driven by genetic factors.

are quite rare, and some pre-parental-death coefficients are statistically significant, the results in Tables A3a–A3d should be interpreted as suggestive, rather than conclusive.

Our second model is a difference-in-differences event study, where we compare individuals with a parental death to a control group of individuals without a parental death, using the event study framework. Figure 2 illustrates the results from these models for males and females, separately. For each figure, the top panel compares individuals with a paternal death to individuals of the same gender without a paternal death, whereas the bottom panel is for maternal deaths. Each line is the coefficient from a separate event study regression, unlike a more traditional difference-in-differences model with an interaction between the two ‘differences’. Tables A4a–A4b report the regression results for males and females, respectively. Again, the reported effects are the change in the hospitalization rates relative to the year before parental death.

The results in Figure 2 display the same basic pattern as the single difference event study models in Figure 1. Males who experience a paternal death have a drastic increase in the likelihood of hospitalization in the year of the death and the following three years, and this difference is statistically different from the hospitalization pattern for males who did not experience a paternal death. Otherwise, any increase in hospitalization due to parental death – maternal or paternal – is indistinguishable from the control group.

Figure 2: Difference-in-Differences Event Study Results for Males and Females



Notes: The figures plot the coefficient estimates from difference-in-differences event study regressions, together with 95% confidence intervals (standard errors clustered at the individual level). The outcome is the hospitalization rate.

With respect to females, we can never distinguish the difference in hospitalization between those experiencing a paternal death and others, similar to previous models. We detect a noticeably higher likelihood of hospitalization in the year of a maternal death and the following year. However, by year 2, the standard error of the treatment group coefficient is so wide that we cannot rule out the possibility (at the 95-percent confidence interval) that the likelihood of hospitalization is the same for the treatment and control groups. This finding, coupled with a similar result for 4 or more years after experiencing a paternal death for males, is the only instance where the event study (Figure 1) and the difference-in-differences event study (Figure 2) differ notably.

In brief, both event study specifications show that individuals who experience the death of a parent of the same gender have dramatically higher hospitalization rates, nearly equal to the mean hospitalization rate for the treatment group, in the year of the parental

death and the following year. The effect persists slightly longer for males, but by four years after the parental death, we cannot detect any significant differences in the coefficients between the treatment and control groups.¹⁵


Next, we estimate event study specifications separately for uninformative and informative causes of death (Figure 3), based on the technique in Gimenez et al. (2013).¹⁶ Because uninformative causes of death, by construction, are unrelated to parental characteristics, the parallel trends assumption should hold. These results reveal two key findings. First, we observe that our baseline conclusions remain intact using only uninformative causes of death that are likely driven by random causes.¹⁷ Second, we find that the effects on hospitalization tend to be quantitatively larger using uninformative causes of death, *vis-a-vis* informative causes of death. This finding is plausible, because children are likely to face greater immediate shock after the incidence of uninformative causes of death that are independent from the socioeconomic characteristics of the family.

¹⁵ These conclusions also hold when we use the imputation estimator for the treatment and control groups (Figures B1 and B2).

¹⁶ See Appendix C for the empirical results on the classification of deaths into UCODs and ICODs.

¹⁷ For females, the effect of maternal death is positive, but no longer significant due to low number of maternal UCODs.

Figure 3: Event Study Coefficients for Hospitalization, ICOD vs. UCOD of a Parent, Males and Females



Notes: The figures plot the coefficient estimates from event study regressions. The outcome is the hospitalization rate. Models were run separately for informative and uninformative parental deaths (ICOD vs UCOD). For clarity, only 95% confidence intervals are shown for the UCODs (standard errors clustered at the individual level).

Heterogeneity by Age Group and Parental Income/education

In our main specifications we use the age range from 10 to 20 due to statistical power reasons given the low parental death rates at earlier ages (Table 1) and control for age fixed effects to account for age-specific vulnerability to mental health problems. Thus, the approach estimates the average impact of parental loss across a rather broad age range of the child. To address this issue further, we investigate the potential heterogeneity of the effects by child age and present the effects between ages 10 and 15 compared to ages 16 to 20. The two main findings, shown in Figure AX and Tables A5a-A5b, hold for both age groups: (1) larger effects for a parental death of a different gender and (2) the longer duration of the effect for a paternal effect. Given that we cannot reject the hypothesis that the coefficient for


ages 10 to 15 is equal to the coefficient for ages 16 to 20, we maintain that pooling the results for the two age groups is appropriate.

We have also estimated the event study models by parental income. This constitutes an approach to isolate the impact of the loss of parental resources. We did not find any noticeable differences in the results by parental income or education (Figure A4 and Tables A6a–A6b).


Family Fixed Effects Results

Next, we proceed to the specifications that incorporate family fixed effects. This model holds constant all time-invariant parental characteristics like mental health and occupational status, instead exploiting within-family differences in the children’s ages at the time of parental death. Figures XX and Table AX show, once again, that the largest negative short-run effects on mental health prevail for those children (boys and girls) who lose a parent of a different gender.

Figure 4: Event Study Results by Age Group at Parental Death




Panel B: Parental death at ages 16-20



Notes: The figures plot the coefficient estimates from event study regressions, together with 95% confidence intervals (standard errors clustered at the individual level). The outcome is the hospitalization rate. Panels on the left show estimates for father's death and those on the right show them for mother's death.

Figure A5: Event Study Results, Family Fixed Effects Regressions



Notes: The figures plot the coefficient estimates from event study regressions, together with 95% confidence intervals (standard errors clustered at the individual level). The outcome is the hospitalization rate. Panels on the left show estimates for father's death and those on the right show them for mother's death.

Cross-Sectional Estimates

Finally, we turn to results from the cross-sectional model estimated using equation (2) for the pooled sample of individuals with and without a parental death. This model provides a longer-term perspective on the association of outcomes with parental death, complementing the in-depth analysis of the short-term outcomes in the event study analysis. The results are based on a linear probability model, where we account for age and year fixed effects, along with extensive demographic control variables (see equation 2).

We summarize the results only very briefly in the main text and describe the findings more fully in Appendix D. We find significant increases in the use of mental health-related medications and sickness absence that are consistent with the increase in the probability of mental-health related hospitalizations, as well as significant reductions in years of schooling, employment, and earnings for the affected children in adulthood. The results for economic consequences of parental death in adulthood echo the findings of the earlier literature.

Although informative, the cross-sectional results – particularly for death at later ages – are best viewed as descriptive for two reasons. First, the models control only for non-random variation in parental death through the inclusion of control variables (X_i). In this model, for example, the hospitalization outcomes could have taken place before the parental death and therefore not be causal. Second, we cannot distinguish whether the smaller coefficients for later parental deaths are because the impact of the death is smaller at later ages or because the individual has fewer years after the parental death to be hospitalized, although both explanations are plausible.

Conclusions

Parental death is a traumatic life event which has major impacts on many life domains. We provide evidence on the effect of parental death on mental health outcomes, with a focus on mental health-related hospitalization. Given the empirical literature in other disciplines on the adverse effects of parental death on mental health, our analysis is a starting point for economists.

Using nationwide register-based data, our results extend the empirical literature in several ways. We find robust evidence that parental death has the most adverse outcomes when males experience a paternal death. The likelihood of hospitalization for mental health reasons roughly doubles – from a very low base of less than 0.01 – in the year of death, and the effect is large and significant in the following 2–3 years (depending on the model). Females experiencing a maternal death also have a near doubling of the (very low) likelihood of hospitalization in the year of death and the following year. In contrast, we generally cannot reject the possibility of no change in the hospitalization rates for males experiencing maternal death or females experiencing paternal death.

The main findings are robust across multiple econometric techniques. Our preferred event study method illustrates the short-run and medium-run outcomes year-by-year. Although the model requires substantial assumptions about exogeneity, these concerns are mitigated by the similarity of results across several dimensions: (1) between models with and without control variables, (2) with and without a control group of individuals who did not experience a parental death, (3) an imputation estimator to allow for heterogeneous treatment effects (Borusyak et al., 2021), (4) to allow restricted to plausibly exogenous causes of death that are not correlated with parental socioeconomic characteristics, as in Espinosa and Evans (2008) and Gimenez et al. (2013), and (5) family fixed effects to account for time-invariant differences between families. Although each technique has its own limitations and assumptions, the consistency across these methods supports a causal effect. Because parental

death is a rare event, and the population of Finland – and other northern European countries with detailed registry data – is moderate, most subgroup analyses are imprecise.

In general, our results are broadly consistent with the economics literature on parental death, the focus of which is on educational and labor-market outcomes, tending to find stronger effects for maternal death rather than paternal death. Our results for females experiencing a maternal death show a consistency between mental health outcomes and educational and labor-market outcomes. Although the economics literature does not find strong effects for paternal death, even among males, for educational and labor-market outcomes, past work has found a relationship between paternal death and other outcomes such as marriage (Lang and Zargosky, 2001) and psychological profile and health (Adda, Björklund, and Holmlund, 2011). Taken collectively, the literature suggests that parental death has a heterogeneous effect on a diverse set of economic outcomes.

Our findings can be used to provide practical guidance for policy setting. Since parental death leads to substantial negative effects on mental health and to significant labor market losses, policy makers should pay special attention to the allocation of appropriate resources to interventions aimed at mitigating these effects. Many of the interventions (e.g., provision of effective mental health services like therapy) would need to occur in schools. Finland has a universal health care system that covers all citizens, but there were no formal nationwide procedures for providing mental health services such as therapy for those children who lost a parent during the time span of our study. The finding that the death of the father has a particularly negative effect on mental health of children may guide income support programs that compensate for the lost family income.

Deeper understanding of the exact mechanisms through which parental death affects mental health, as well as the heterogeneous effects of parental death, are important topics for future empirical research. Although our results are consistent with mechanisms identified in the broader social science, we cannot identify the direct mechanisms at play because the

register data are not well suitable for the examination of social interactions (including social support) and the degree of emotional attachment within families. Identifying the exact mechanisms is also hampered due to the lack of suitable survey data and limited earlier evidence on social connections in the family using Finnish data.

References

- Adda, Jérôme, Anders Björklund, and Helena Holmlund. 2011. The Role of Mothers and Fathers in Providing Skills: Evidence from Parental Deaths. IZA Discussion Paper Number 5425.
- Alexander, Diane, and Molly Schnell. 2019. Just What the Nurse Practitioner Ordered: Independent Prescriptive Authority and Population Mental Health. *Journal of Health Economics* 66: 145–162.
- Altonji, Joseph G., Todd E. Elder, and Christopher R. Taber. 2005. Selection on Observed and Unobserved Variables: Assessing the Effectiveness of Catholic School. *Journal of Political Economy* 113(1): 151–184.
- Amato, Paul R. 1993. Children's Adjustment to Divorce: Theories, Hypotheses, and Empirical Support. *Journal of Marriage and the Family*: 55(1): 23–38.
- Appel, Charlotte Weiling, Christoffer Johansen, Isabelle Deltour, Kirsten Frederiksen, Henrik Hjalgrim, Susanne Oksbjerg Dalton, Annemarie Dencker, JesDige, Per Bøge, Bo Andreassen Rix, Atle Dyregrov, PrebenEngelbrekt, Eva Helweg, Ole Abildgaard Mikkelsen, Mette TerpHøybye, and Pernille Envold Bidstrup. 2013. Early Parental Death and Risk of Hospitalization for Affective Disorder in Adulthood. *Epidemiology* 24(4): 608–661.
- Berg, Lisa, Mikael Rostila, and Anders Hjern. 2016. Parental Death during Childhood and Depression in Young Adults – A National Cohort Study. *The Journal of Child Psychology and Psychiatry* 57(9): 1092–1098.
- Bokhorst, Caroline L., Sindy R. Sumter and P. Michiel Westenberg. 2010. Social Support from Parents, Friends, Classmates, and Teachers in Children and Adolescents Aged 9 to 18 Years: Who is Perceived as Most Supportive? *Social Development* 19(2): 417–426.
- Borusyak, K., Xavier Jaravel, and Jann Spiess. 2021. Revisiting Event Study Designs: Robust and Efficient Estimation. Mimeo.
- Bubonya, Melisa, Deborah Cobb-Clark, and Mark Wooden. 2017. Mental Health and Productivity at Work: Does What You Do Matter? *Labour Economics* 46: 150–165.
- Böckerman, Petri, Ohto Kanninen, and Ippo Suoniemi. 2018. A Kink that Makes You Sick: The Effect of Sick Pay on Absence. *Journal of Applied Econometrics* 33: 568–579.

- Chen, Stacey H., Yen-Chien Chen, and Jin-Tan Liu. 2009. The Impact of Unexpected Maternal Death on Education: First Evidence from Three National Administrative Data Links. *American Economic Review: Papers and Proceedings* 99(2): 149–153.
- Corak, Miles. 2001. Death and Divorce: The Long-Term Consequences of Parental Loss on Adolescents. *Journal of Labor Economics* 19(3): 682–715.
- Dupraz, Yannick, and Andy Ferrara. 2021. Fatherless: The Long-Term Effects of Losing a Father in the U.S. Civil War. CAGE Working Paper no. 538.
- Espinosa, Javier, and William N. Evans. 2008. Heightened Mortality after the Death of a Spouse: Marriage Protection or Marriage Selection? *Journal of Health Economics* 27 (5): 1326–1342.
- Freyaldenhoven, S., Hansen, C., Pérez, J.P., Shapiro, J.M. (2021). Visualization, Identification, and Estimation in the Linear Panel Event-Study Design. National Bureau of Economic Research, Working Paper No. 29170.
- Fronstin, Paul, David H. Greenberg, and Philip K. Robins. 2001. Parental Disruption and the Labour Market Performance of Children When They Reach Adulthood. *Journal of Population Economics* 14(1): 137–172.
- Gimenez, Lea, Shin-Yi Chou, Jin-Tan Liu, and Jin-Long Liu. 2013. Parental Loss and Children's Well Being. *Journal of Human Resources* 48(4): 1035–1071
- Gould, Eric, Avi Simhon, and Bruce A. Weinberg. 2019. Does Parental Quality Matter? Evidence on the Transmission of Human Capital Using Variation in Parental Influence from Death, Divorce, and Family Size. National Bureau of Economic Research Working Paper Number 25495.
- Hakulinen, C., Elovainio, M., Arffman, M., Lumme, S., Pirkola, S., Keskimäki, I., Manderbacka, P. and Böckerman, P. 2019. Mental Disorders and Long-term Labour Market Outcomes: Nationwide Cohort Study of 2 055 720 Individuals. *Acta Psychiatrica Scandinavica* 140(4): 371–381.
- Kailaheimo-Lönnqvist, Sanna, and Jani Erola. 2020. Child's Age at Parental Death and University Education. *European Societies* 22(4) 433–455.
- Kalil, Ariel, Magne Mogstad, Mari Rege, and Mark E. Votruba. 2016. Father Presence and the Intergenerational Transmission of Educational Attainment. *Journal of Human Resources* 51(4): 869–899.
- Kleven, Henrik, Camille Landais, and Jakob Egholt Søgaard. 2019. Children and Gender Inequality: Evidence from Denmark. *American Economic Journal: Applied Economics* 11(4): 181–209.
- Kristiansen, Ida Lykke. 2021. Consequences of Serious Parental Health Events on Child Mental Health and Educational Outcomes. *Health Economics*, Forthcoming. Doi: <https://doi.org/10.1002/hec.4278>
- Lahti, Raimo A., and Antti Penttilä. 2001. The Validity of Death Certificates: Routine Validation of Death Certification and Its Effects on Mortality Statistics. *Forensic Science International* 115(1–2): 15–32.

- Lang, Kevin, and Jay L. Zagorsky. 2001. Does Growing Up with a Parent Absent Really Hurt? *Journal of Human Resources* 36(2): 253–273.
- Layard, Richard. 2013. Mental Health: The New Frontier for Labour Economics. *IZA Journal of Labor Policy* 2: 1–16.
- Liu, Nancy H., Daumit, Gail L., Dua, Tarun, ... and Saxena, Shekhar. 2017. Excess Mortality in Persons with Severe Mental Disorders: A Multilevel Intervention Framework and Priorities for Clinical Practice, Policy and Research Agendas. *World Psychiatry* 16: 30–40.
- McKay, Michael T., Mary Cannon, Colm Healy, Sarah Syer, Laurie O’Donnell, and Mary C. Clarke. 2021. A Meta-Analysis of the Relationship between Parental Death in Childhood and Subsequent Psychiatric Disorder. *Acta Psychiatrica Scandinavica* Forthcoming.
- Persson, Petra, and Maya Rossin-Slater. 2018. Family Ruptures, Stress, and the Mental Health of the Next Generation. *American Economic Review* 108(4–5): 1214–1252.
- Raley, Sara, and Suzanne Bianchi. 2006. Sons, Daughters, and Family Processes: Does Gender of Children Matter? *Annual Review of Sociology* 32: 401–421.
- Rosenbaum-Feldbrügge, Matthias. 2019. The Impact of Parental Death in Childhood on Sons’ and Daughters’ Status Attainment in Young Adulthood in the Netherlands, 1850-1952. *Demography* 56(5): 1827–1854.
- Räikkönen, Katri, Marius Lahti, Kati Heinonen, Anu-Katriina Pesonen, Kristian Wahlbeck, Eero Kajantie, ... and Johan G. Eriksson. (2011). Risk of severe mental disorders in adults separated temporarily from their parents in childhood: the Helsinki birth cohort study. *Journal of Psychiatric Research* 45: 332-338.
- Saarela, Jan and Fjalar Finnäs. 2014. Sex Composition of Children, Parental Separation, and Parity Progression: Is Finland a Nordic Outlier? *Demographic Research* 30: 49–70.
- Sareen, Jitender, Frank Jacobi, Brian J. Cox, Shay-Lee Belik, Ian Clara, and Murray B. Stein. 2006. Disability and Poor Quality of Life Associated with Comorbid Anxiety Disorders and Physical Conditions. *Archives of Internal Medicine* 166: 2109–2116.
- Steele, Fiona, Wendy Sigle-Rushton, and Øystein Kravdal. 2009. Consequences of Family Disruption on Children’s Educational Outcomes in Norway. *Demography* 46(3): 553–574.
- Sund, Reijo. 2012. Quality of the Finnish Hospital Discharge Register: A Systematic Review. *Scandinavian Journal of Public Health* 40: 505–515.
- Tousoulis, Dimitris, Evangelos Oikonomou, Georgia Vogiatzi, and Panos Vardas. 2020. Cardiovascular Disease and Socioeconomic Status: It Is Mainly Education that Counts and not Wealth! *European Heart Journal* 41: 3213–3214.
- von Bismarck-Osten, Clara, Kirill Borusyak and Uta Schönberg. 2020. The Role of Schools in Transmission of the SARS-CoV-2 Virus: Quasi-Experimental Evidence from Germany. CReAM Discussion Paper Series 2022, Centre for Research and Analysis of Migration (CReAM), University College London.

Whiteford, Harvey A, Louisa Degenhardt, Jürgen Rehm, et al. 2013. Global Burden of Disease Attributable to Mental and Substance Use Disorders: Findings from the Global Burden of Disease Study 2010. *The Lancet* 382: 1575–1586.

Wiese, Bettina S., and Alexandra M. Freund. 2011. Parents as Role Models: Parental Behavior Affects Adolescents' Plans for Work Involvement. *International Journal of Behavioral Development* 35: 218–224.

SUPPLEMENTARY ONLINE APPENDIX (NOT FOR PRINT)

CONTENT

- Appendix A. Tables and Figures: Event Study Design
- Appendix B. Pre-Trend and Imputation Estimation Results
- Appendix C. Uninformative vs. Informative Parental Deaths
- Appendix D. Tables and Figures: Cross-Sectional Analysis

APPENDIX A. TABLES AND FIGURES: EVENT STUDY DESIGN

Table A1: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|--------------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | 0.0003 (0.0008) | -0.0006 (0.0015) | 0.0004 (0.0006) | -0.0003 (0.0012) |
| 7 years before death | 0.0008 (0.0008) | 0.0017 (0.0015) | 0.0006 (0.0006) | 0.0010 (0.0012) |
| 6 years before death | 0.0008 (0.0007) | -0.0006 (0.0014) | 0.0007 (0.0006) | -0.0002 (0.0011) |
| 5 years before death | 0.0009 (0.0007) | 0.0002 (0.0013) | 0.0003 (0.0006) | 0.0003 (0.0011) |
| 4 years before death | 0.0006 (0.0007) | -0.0003 (0.0012) | 0.0003 (0.0006) | 0.0005 (0.0011) |
| 3 years before death | 0.0003 (0.0006) | 0.0018 (0.0013) | -0.0003 (0.0006) | 0.0013 (0.0011) |
| 2 years before death | 0.0006 (0.0006) | 0.0006 (0.0011) | 0.0005 (0.0006) | 0.0013 (0.0010) |
| Year of death | 0.0044*** (0.0008) | 0.0028* (0.0015) | 0.0004 (0.0006) | 0.0039*** (0.0011) |
| 1 year after death | 0.0039*** (0.0009) | 0.0020 (0.0015) | 0.0004 (0.0007) | 0.0033*** (0.0013) |
| 2 years after death | 0.0029*** (0.0009) | 0.0001 (0.0015) | 0.0014* (0.0008) | 0.0028** (0.0014) |
| 3 years after death | 0.0029*** (0.0010) | 0.0002 (0.0016) | 0.0012 (0.0008) | 0.0018 (0.0014) |
| 4 years after death | 0.0014 (0.0010) | 0.0003 (0.0017) | 0.0012 (0.0009) | 0.0006 (0.0015) |
| 5 years after death | 0.0021** (0.0010) | -0.0008 (0.0018) | 0.0007 (0.0010) | 0.0008 (0.0016) |
| 6 years after death | 0.0025** (0.0011) | 0.0012 (0.0021) | 0.0006 (0.0010) | 0.0009 (0.0018) |
| 7 years after death | 0.0023* (0.0012) | 0.0006 (0.0022) | 0.0005 (0.0011) | 0.0004 (0.0019) |
| 8 years after death | 0.0017 (0.0014) | 0.0030 (0.0024) | 0.0002 (0.0012) | 0.0009 (0.0021) |
| 9 years after death | 0.0025* (0.0015) | 0.0012 (0.0027) | 0.0008 (0.0012) | 0.0008 (0.0023) |
| 10 years after death | 0.0035** (0.0017) | 0.0004 (0.0030) | 0.0002 (0.0014) | -0.0002 (0.0025) |
| Observations | 343,600 | 125,187 | 334,878 | 118,081 |
| Number of individuals | 18,272 | 6,676 | 17,798 | 6,299 |
| R-squared | 0.0038 | 0.0044 | 0.0028 | 0.0037 |
| Age fixed effects | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES |
| Individual fixed effects | NO | NO | NO | NO |
| Additional controls | NO | NO | NO | NO |
| Mean $Y_{t=1}$ | 0.0045 | 0.0062 | 0.0039 | 0.0037 |

Notes: Standard errors clustered at the individual level in parentheses. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20.

Table A2a: Alternate Event Study Models, By Parental Death, Males

| | Father (1) | Father (2) | Father (3) | Mother (4) | Mother (5) | Mother (6) |
|--------------------------|-----------------------|-----------------------|-----------------------|---------------------|----------------------|---------------------|
| 8 years before death | 0.0003 (0.0008) | -0.0015 (0.0016) | 0.0007 (0.0008) | -0.0006 (0.0015) | -0.0032 (0.0028) | 0.0006 (0.0015) |
| 7 years before death | 0.0008 (0.0008) | -0.0008 (0.0015) | 0.0012 (0.0008) | 0.0017 (0.0015) | -0.0005 (0.0025) | 0.0027* (0.0015) |
| 6 years before death | 0.0008 (0.0007) | -0.0004 (0.0013) | 0.0012* (0.0007) | -0.0006 (0.0014) | -0.0023 (0.0023) | 0.0002 (0.0014) |
| 5 years before death | 0.0009 (0.0007) | -0.0001 (0.0011) | 0.0012* (0.0007) | 0.0002 (0.0013) | -0.0011 (0.0019) | 0.0009 (0.0013) |
| 4 years before death | 0.0006 (0.0007) | -0.0002 (0.0010) | 0.0006 (0.0007) | -0.0003 (0.0012) | -0.0013 (0.0016) | 0.0002 (0.0012) |
| 3 years before death | 0.0003 (0.0006) | -0.0002 (0.0008) | 0.0003 (0.0006) | 0.0018 (0.0013) | 0.0011 (0.0015) | 0.0021 (0.0013) |
| 2 years before death | 0.0006 (0.0006) | 0.0003 (0.0007) | 0.0006 (0.0006) | 0.0006 (0.0011) | 0.0002 (0.0012) | 0.0007 (0.0011) |
| Year of death | 0.0044*** (0.0008) | 0.0047*** (0.0009) | 0.0042*** (0.0008) | 0.0028* (0.0015) | 0.0031** (0.0016) | 0.0026* (0.0015) |
| 1 year after death | 0.0039*** (0.0009) | 0.0045*** (0.0010) | 0.0036*** (0.0009) | 0.0020 (0.0015) | 0.0026 (0.0017) | 0.0016 (0.0015) |
| 2 years after death | 0.0029*** (0.0009) | 0.0038*** (0.0011) | 0.0024*** (0.0009) | 0.0001 (0.0015) | 0.0012 (0.0020) | -0.0004 (0.0015) |
| 3 years after death | 0.0029*** (0.0010) | 0.0040*** (0.0012) | 0.0026*** (0.0010) | 0.0002 (0.0016) | 0.0017 (0.0023) | -0.0004 (0.0016) |
| 4 years after death | 0.0014 (0.0010) | 0.0028** (0.0014) | 0.0012 (0.0010) | 0.0003 (0.0017) | 0.0021 (0.0025) | -0.0005 (0.0017) |
| 5 years after death | 0.0021** (0.0010) | 0.0037** (0.0015) | 0.0016 (0.0010) | -0.0008 (0.0018) | 0.0013 (0.0029) | -0.0019 (0.0018) |
| 6 years after death | 0.0025** (0.0011) | 0.0044** (0.0017) | 0.0018* (0.0011) | 0.0012 (0.0021) | 0.0036 (0.0032) | 0.0000 (0.0021) |
| 7 years after death | 0.0023* (0.0012) | 0.0045** (0.0019) | 0.0021* (0.0012) | 0.0006 (0.0022) | 0.0032 (0.0037) | -0.0008 (0.0022) |
| 8 years after death | 0.0017 (0.0014) | 0.0041** (0.0020) | 0.0016 (0.0013) | 0.0030 (0.0024) | 0.0058 (0.0041) | 0.0015 (0.0025) |
| 9 years after death | 0.0025* (0.0015) | 0.0054** (0.0023) | 0.0022 (0.0015) | 0.0012 (0.0027) | 0.0043 (0.0045) | -0.0005 (0.0028) |
| 10 years after death | 0.0035** (0.0017) | 0.0066*** (0.0023) | 0.0027 (0.0017) | 0.0004 (0.0030) | 0.0035 (0.0048) | -0.0015 (0.0030) |
| Observations | 343,600 | 343,600 | 330,905 | 125,187 | 125,187 | 125,187 |
| No of individuals | 18,272 | 18,272 | 17,485 | 6,676 | 6,676 | 6,676 |
| R-squared | 0.0038 | 0.1480 | 0.0066 | 0.0044 | 0.1637 | 0.0113 |
| Age fixed effects | YES | YES | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES | YES | YES |
| Individual fixed effects | NO | YES | NO | NO | YES | NO |
| Additional controls | NO | NO | YES | NO | NO | YES |
| Mean Y_{t-1} | 0.0045 | 0.0045 | 0.0042 | 0.0062 | 0.0062 | 0.0062 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20.

Table A2b: Alternate Event Study Models, By Parental Death, Females

| | Father (1) | Father (2) | Father (3) | Mother (4) | Mother (5) | Mother (6) |
|--------------------------|---------------------|----------------------|---------------------|-----------------------|-----------------------|-----------------------|
| 8 years before death | 0.0004 (0.0006) | -0.0025 (0.0018) | 0.0008 (0.0006) | -0.0003 (0.0012) | -0.0014 (0.0013) | 0.0000 (0.0012) |
| 7 years before death | 0.0006 (0.0006) | -0.0018 (0.0016) | 0.0010 (0.0006) | 0.0010 (0.0012) | -0.0001 (0.0014) | 0.0013 (0.0012) |
| 6 years before death | 0.0007 (0.0006) | -0.0013 (0.0013) | 0.0010 (0.0006) | -0.0002 (0.0011) | -0.0011 (0.0012) | 0.0000 (0.0011) |
| 5 years before death | 0.0003 (0.0006) | -0.0013 (0.0011) | 0.0005 (0.0006) | 0.0003 (0.0011) | -0.0004 (0.0012) | 0.0005 (0.0011) |
| 4 years before death | 0.0003 (0.0006) | -0.0009 (0.0010) | 0.0005 (0.0006) | 0.0005 (0.0011) | -0.0000 (0.0011) | 0.0007 (0.0011) |
| 3 years before death | -0.0003 (0.0006) | -0.0011 (0.0008) | -0.0002 (0.0006) | 0.0013 (0.0011) | 0.0009 (0.0011) | 0.0014 (0.0011) |
| 2 years before death | 0.0005 (0.0006) | 0.0001 (0.0007) | 0.0007 (0.0006) | 0.0013 (0.0010) | 0.0011 (0.0010) | 0.0013 (0.0010) |
| Year of death | 0.0004 (0.0006) | 0.0008 (0.0007) | 0.0003 (0.0006) | 0.0039*** (0.0011) | 0.0040*** (0.0012) | 0.0038*** (0.0011) |
| 1 year after death | 0.0004 (0.0007) | 0.0012 (0.0009) | 0.0005 (0.0007) | 0.0033*** (0.0013) | 0.0036*** (0.0013) | 0.0032** (0.0013) |
| 2 years after death | 0.0014* (0.0008) | 0.0026** (0.0010) | 0.0012 (0.0008) | 0.0028** (0.0014) | 0.0033** (0.0014) | 0.0027** (0.0014) |
| 3 years after death | 0.0012 (0.0008) | 0.0028** (0.0012) | 0.0011 (0.0008) | 0.0018 (0.0014) | 0.0025* (0.0014) | 0.0016 (0.0014) |
| 4 years after death | 0.0012 (0.0009) | 0.0031** (0.0014) | 0.0007 (0.0009) | 0.0006 (0.0015) | 0.0015 (0.0015) | 0.0004 (0.0015) |
| 5 years after death | 0.0007 (0.0010) | 0.0030* (0.0016) | 0.0006 (0.0010) | 0.0008 (0.0016) | 0.0020 (0.0015) | 0.0006 (0.0016) |
| 6 years after death | 0.0006 (0.0010) | 0.0034* (0.0019) | 0.0006 (0.0010) | 0.0009 (0.0018) | 0.0022 (0.0016) | 0.0006 (0.0018) |
| 7 years after death | 0.0005 (0.0011) | 0.0036* (0.0021) | 0.0001 (0.0011) | 0.0004 (0.0019) | 0.0019 (0.0018) | 0.0001 (0.0019) |
| 8 years after death | 0.0002 (0.0012) | 0.0036 (0.0023) | -0.0001 (0.0012) | 0.0009 (0.0021) | 0.0025 (0.0019) | 0.0006 (0.0021) |
| 9 years after death | 0.0008 (0.0012) | 0.0045* (0.0026) | 0.0006 (0.0013) | 0.0008 (0.0023) | 0.0025 (0.0020) | 0.0004 (0.0023) |
| 10 years after death | 0.0002 (0.0014) | 0.0042 (0.0027) | 0.0001 (0.0014) | -0.0002 (0.0025) | 0.0018 (0.0018) | -0.0006 (0.0025) |
| Observations | 334,878 | 334,878 | 324,440 | 118,081 | 118,081 | 118,081 |
| Number of individuals | 17,798 | 17,798 | 17,157 | 6,299 | 6,299 | 6,299 |
| R-squared | 0.0028 | 0.1599 | 0.0053 | 0.0037 | 0.1710 | 0.0082 |
| Age fixed effects | YES | YES | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES | YES | YES |
| Individual fixed effects | NO | YES | NO | NO | YES | NO |
| Additional controls | NO | NO | YES | NO | NO | YES |
| Mean Y_{t-1} | 0.0039 | 0.0039 | 0.0039 | 0.0037 | 0.0037 | 0.0037 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20.

Table A3a: Event Study Models by Cause of Hospitalization, Paternal Death, Males

| | All (1) | Depression (2) | Anxiety (3) | Substance (4) | Self-harm (5) |
|----------------------|-----------------------|-----------------------|----------------------|---------------------|----------------------|
| 8 years before death | 0.0003 (0.0008) | 0.0001 (0.0002) | -0.0001 (0.0002) | -0.0000 (0.0002) | 0.0001 (0.0001) |
| 7 years before death | 0.0008 (0.0008) | 0.0000 (0.0002) | -0.0000 (0.0002) | 0.0000 (0.0002) | 0.0001 (0.0001) |
| 6 years before death | 0.0008 (0.0007) | -0.0000 (0.0002) | 0.0001 (0.0002) | 0.0000 (0.0002) | 0.0000 (0.0001) |
| 5 years before death | 0.0009 (0.0007) | 0.0001 (0.0002) | -0.0000 (0.0002) | -0.0001 (0.0002) | 0.0001 (0.0001) |
| 4 years before death | 0.0006 (0.0007) | 0.0003 (0.0003) | 0.0000 (0.0001) | -0.0001 (0.0002) | 0.0002 (0.0002) |
| 3 years before death | 0.0003 (0.0006) | 0.0000 (0.0002) | 0.0002 (0.0002) | 0.0001 (0.0003) | 0.0001 (0.0001) |
| 2 years before death | 0.0006 (0.0006) | -0.0000 (0.0002) | 0.0003 (0.0002) | -0.0002 (0.0003) | 0.0001 (0.0002) |
| Year of death | 0.0044*** (0.0008) | 0.0008** (0.0003) | 0.0005** (0.0002) | 0.0005 (0.0003) | 0.0002 (0.0002) |
| 1 year after death | 0.0039*** (0.0009) | 0.0010*** (0.0003) | 0.0002 (0.0002) | 0.0008* (0.0004) | 0.0002 (0.0002) |
| 2 years after death | 0.0029*** (0.0009) | 0.0009** (0.0004) | 0.0001 (0.0002) | 0.0004 (0.0004) | 0.0005** (0.0002) |
| 3 years after death | 0.0029*** (0.0010) | 0.0005 (0.0003) | 0.0004 (0.0003) | -0.0000 (0.0004) | 0.0004 (0.0003) |
| 4 years after death | 0.0014 (0.0010) | -0.0001 (0.0003) | 0.0003 (0.0002) | -0.0004 (0.0005) | 0.0002 (0.0002) |
| 5 years after death | 0.0021** (0.0010) | 0.0003 (0.0004) | 0.0004 (0.0003) | 0.0002 (0.0005) | 0.0006** (0.0003) |
| 6 years after death | 0.0025** (0.0011) | 0.0001 (0.0003) | 0.0006** (0.0003) | 0.0001 (0.0006) | 0.0004 (0.0003) |
| 7 years after death | 0.0023* (0.0012) | 0.0007* (0.0004) | 0.0005* (0.0003) | -0.0000 (0.0006) | 0.0004 (0.0003) |
| 8 years after death | 0.0017 (0.0014) | -0.0002 (0.0004) | 0.0008** (0.0003) | 0.0002 (0.0007) | 0.0006* (0.0004) |
| 9 years after death | 0.0025* (0.0015) | 0.0002 (0.0005) | 0.0007** (0.0003) | 0.0001 (0.0008) | 0.0007* (0.0004) |
| 10 years after death | 0.0035** (0.0017) | 0.0003 (0.0005) | 0.0007* (0.0004) | 0.0005 (0.0009) | 0.0006 (0.0004) |
| Observations | 343,600 | 343,600 | 343,600 | 343,600 | 343,600 |
| No of individuals | 18,272 | 18,272 | 18,272 | 18,272 | 18,272 |
| R-squared | 0.0038 | 0.0009 | 0.0010 | 0.0023 | 0.0008 |
| Mean Y_{t-1} | 0.0045 | 0.00049 | 0.00022 | 0.00088 | 0.00022 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20. All models contain year and age fixed effects.

Table A3b: Event Study Models by Cause of Hospitalization, Maternal Death, Males

| | All (1) | Depression (2) | Anxiety (3) | Substance (4) | Self-harm (5) |
|----------------------|---------------------|---------------------|---------------------|----------------------|---------------------|
| 8 years before death | -0.0006 (0.0015) | 0.0004 (0.0005) | 0.0003* (0.0002) | 0.0003 (0.0004) | -0.0000 (0.0002) |
| 7 years before death | 0.0017 (0.0015) | 0.0003 (0.0005) | 0.0004* (0.0002) | 0.0001 (0.0004) | -0.0001 (0.0002) |
| 6 years before death | -0.0006 (0.0014) | -0.0001 (0.0004) | 0.0003* (0.0002) | 0.0001 (0.0004) | -0.0001 (0.0002) |
| 5 years before death | 0.0002 (0.0013) | -0.0003 (0.0004) | 0.0004* (0.0002) | 0.0001 (0.0004) | -0.0002 (0.0002) |
| 4 years before death | -0.0003 (0.0012) | -0.0004 (0.0004) | 0.0003* (0.0002) | 0.0001 (0.0004) | 0.0001 (0.0003) |
| 3 years before death | 0.0018 (0.0013) | -0.0004 (0.0004) | 0.0002 (0.0002) | 0.0004 (0.0004) | -0.0001 (0.0003) |
| 2 years before death | 0.0006 (0.0011) | -0.0003 (0.0004) | 0.0005* (0.0003) | 0.0004 (0.0005) | 0.0003 (0.0003) |
| Year of death | 0.0028* (0.0015) | 0.0000 (0.0006) | 0.0008* (0.0004) | 0.0003 (0.0006) | 0.0002 (0.0004) |
| 1 year after death | 0.0020 (0.0015) | 0.0006 (0.0006) | 0.0001 (0.0004) | 0.0005 (0.0006) | 0.0002 (0.0004) |
| 2 years after death | 0.0001 (0.0015) | -0.0008 (0.0005) | -0.0003 (0.0003) | 0.0003 (0.0007) | 0.0004 (0.0004) |
| 3 years after death | 0.0002 (0.0016) | -0.0004 (0.0006) | 0.0001 (0.0004) | 0.0002 (0.0007) | 0.0003 (0.0004) |
| 4 years after death | 0.0003 (0.0017) | -0.0001 (0.0007) | -0.0000 (0.0004) | 0.0005 (0.0008) | -0.0002 (0.0004) |
| 5 years after death | -0.0008 (0.0018) | 0.0003 (0.0007) | -0.0001 (0.0005) | 0.0005 (0.0008) | 0.0003 (0.0005) |
| 6 years after death | 0.0012 (0.0021) | -0.0003 (0.0007) | 0.0010 (0.0006) | 0.0002 (0.0009) | 0.0001 (0.0006) |
| 7 years after death | 0.0006 (0.0022) | 0.0005 (0.0008) | 0.0003 (0.0006) | 0.0007 (0.0011) | 0.0005 (0.0006) |
| 8 years after death | 0.0030 (0.0024) | -0.0002 (0.0008) | 0.0006 (0.0006) | 0.0027** (0.0012) | 0.0000 (0.0006) |
| 9 years after death | 0.0012 (0.0027) | -0.0005 (0.0008) | -0.0003 (0.0005) | 0.0009 (0.0013) | 0.0001 (0.0007) |
| 10 years after death | 0.0004 (0.0030) | 0.0001 (0.0010) | -0.0001 (0.0005) | 0.0005 (0.0014) | -0.0001 (0.0007) |
| Observations | 125,187 | 125,187 | 125,187 | 125,187 | 125,187 |
| No of individuals | 6,676 | 6,676 | 6,676 | 6,676 | 6,676 |
| R-squared | 0.0044 | 0.0012 | 0.0017 | 0.0031 | 0.0014 |
| Mean $Y_{t=1}$ | 0.0062 | 0.00090 | 0.00015 | 0.00075 | 0.00030 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20. All models contain year and age fixed effects.

Table A3c: Event Study Models by Cause of Hospitalization, Paternal Death, Females

| | All (1) | Depression (2) | Anxiety (3) | Substance (4) | Self-harm (5) |
|----------------------|---------------------|---------------------|---------------------|----------------------|-----------------------|
| 8 years before death | 0.0004 (0.0006) | 0.0002 (0.0003) | -0.0000 (0.0001) | -0.0000 (0.0002) | 0.0001 (0.0002) |
| 7 years before death | 0.0006 (0.0006) | 0.0001 (0.0003) | -0.0000 (0.0001) | -0.0000 (0.0002) | 0.0001 (0.0002) |
| 6 years before death | 0.0007 (0.0006) | 0.0001 (0.0003) | 0.0000 (0.0002) | 0.0000 (0.0002) | 0.0000 (0.0002) |
| 5 years before death | 0.0003 (0.0006) | 0.0003 (0.0003) | -0.0001 (0.0001) | 0.0000 (0.0002) | 0.0002 (0.0003) |
| 4 years before death | 0.0003 (0.0006) | 0.0001 (0.0003) | -0.0001 (0.0001) | 0.0001 (0.0002) | 0.0003 (0.0003) |
| 3 years before death | -0.0003 (0.0006) | -0.0001 (0.0003) | 0.0000 (0.0002) | -0.0001 (0.0002) | -0.0000 (0.0003) |
| 2 years before death | 0.0005 (0.0006) | 0.0000 (0.0003) | -0.0001 (0.0001) | 0.0000 (0.0002) | -0.0000 (0.0003) |
| Year of death | 0.0004 (0.0006) | -0.0004 (0.0003) | 0.0002 (0.0002) | -0.0002 (0.0002) | 0.0001 (0.0003) |
| 1 year after death | 0.0004 (0.0007) | 0.0004 (0.0004) | 0.0001 (0.0002) | 0.0003 (0.0003) | -0.0001 (0.0003) |
| 2 years after death | 0.0014* (0.0008) | 0.0007 (0.0005) | 0.0001 (0.0002) | 0.0003 (0.0003) | 0.0000 (0.0004) |
| 3 years after death | 0.0012 (0.0008) | 0.0006 (0.0005) | -0.0000 (0.0002) | -0.0000 (0.0003) | -0.0001 (0.0004) |
| 4 years after death | 0.0012 (0.0009) | 0.0004 (0.0005) | 0.0005* (0.0003) | 0.0008** (0.0004) | -0.0005 (0.0004) |
| 5 years after death | 0.0007 (0.0010) | 0.0000 (0.0005) | 0.0001 (0.0003) | 0.0002 (0.0004) | -0.0007* (0.0004) |
| 6 years after death | 0.0006 (0.0010) | -0.0000 (0.0005) | 0.0002 (0.0003) | -0.0001 (0.0004) | -0.0006 (0.0004) |
| 7 years after death | 0.0005 (0.0011) | 0.0002 (0.0006) | -0.0000 (0.0003) | 0.0003 (0.0005) | -0.0009** (0.0004) |
| 8 years after death | 0.0002 (0.0012) | 0.0002 (0.0006) | 0.0002 (0.0003) | 0.0000 (0.0005) | -0.0003 (0.0005) |
| 9 years after death | 0.0008 (0.0012) | 0.0008 (0.0006) | 0.0003 (0.0003) | 0.0002 (0.0005) | -0.0007 (0.0005) |
| 10 years after death | 0.0002 (0.0014) | 0.0002 (0.0007) | -0.0001 (0.0003) | 0.0004 (0.0006) | -0.0006 (0.0005) |
| Observations | 334,878 | 334,878 | 334,878 | 334,878 | 334,878 |
| No of individuals | 17,798 | 17,798 | 17,798 | 17,798 | 17,798 |
| R-squared | 0.0028 | 0.0020 | 0.0006 | 0.0012 | 0.0010 |
| Mean Y_{t-1} | 0.00389 | 0.00107 | 0.00028 | 0.00051 | 0.00085 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20. All models contain year and age fixed effects.

Table A3d: Event Study Models by Cause of Hospitalization, Maternal Death, Females

| | All (1) | Depression (2) | Anxiety (3) | Substance (4) | Self-harm (5) |
|----------------------|-----------------------|-----------------------|---------------------|---------------------|-----------------------|
| 8 years before death | -0.0003 (0.0012) | 0.0006 (0.0004) | -0.0001 (0.0002) | 0.0001 (0.0002) | 0.0003 (0.0002) |
| 7 years before death | 0.0010 (0.0012) | 0.0006* (0.0003) | -0.0001 (0.0002) | 0.0001 (0.0002) | 0.0003* (0.0002) |
| 6 years before death | -0.0002 (0.0011) | 0.0007* (0.0004) | -0.0001 (0.0002) | 0.0001 (0.0002) | 0.0002 (0.0002) |
| 5 years before death | 0.0003 (0.0011) | 0.0005 (0.0004) | -0.0001 (0.0002) | 0.0002 (0.0003) | 0.0004 (0.0003) |
| 4 years before death | 0.0005 (0.0011) | 0.0005 (0.0004) | -0.0001 (0.0002) | 0.0000 (0.0003) | 0.0002 (0.0002) |
| 3 years before death | 0.0013 (0.0011) | 0.0007 (0.0005) | -0.0001 (0.0002) | 0.0000 (0.0003) | -0.0001 (0.0002) |
| 2 years before death | 0.0013 (0.0010) | 0.0002 (0.0004) | 0.0001 (0.0002) | 0.0004 (0.0004) | 0.0000 (0.0002) |
| Year of death | 0.0039*** (0.0011) | 0.0011* (0.0006) | 0.0004 (0.0004) | 0.0003 (0.0004) | 0.0013** (0.0005) |
| 1 year after death | 0.0033*** (0.0013) | 0.0021*** (0.0008) | 0.0001 (0.0003) | 0.0006 (0.0004) | 0.0018*** (0.0006) |
| 2 years after death | 0.0028** (0.0014) | 0.0016** (0.0008) | 0.0001 (0.0004) | 0.0004 (0.0004) | 0.0006 (0.0005) |
| 3 years after death | 0.0018 (0.0014) | 0.0013 (0.0009) | 0.0003 (0.0004) | 0.0004 (0.0004) | 0.0007 (0.0005) |
| 4 years after death | 0.0006 (0.0015) | -0.0003 (0.0008) | 0.0007 (0.0005) | -0.0002 (0.0004) | 0.0009 (0.0006) |
| 5 years after death | 0.0008 (0.0016) | -0.0010 (0.0008) | 0.0009* (0.0005) | -0.0001 (0.0006) | 0.0004 (0.0006) |
| 6 years after death | 0.0009 (0.0018) | -0.0003 (0.0009) | -0.0002 (0.0004) | 0.0010 (0.0008) | 0.0003 (0.0006) |
| 7 years after death | 0.0004 (0.0019) | -0.0001 (0.0010) | 0.0000 (0.0005) | 0.0005 (0.0008) | -0.0002 (0.0006) |
| 8 years after death | 0.0009 (0.0021) | -0.0004 (0.0012) | 0.0010 (0.0006) | 0.0002 (0.0008) | 0.0002 (0.0007) |
| 9 years after death | 0.0008 (0.0023) | 0.0008 (0.0013) | -0.0005 (0.0004) | 0.0005 (0.0010) | 0.0001 (0.0007) |
| 10 years after death | -0.0002 (0.0025) | -0.0003 (0.0013) | -0.0001 (0.0005) | 0.0000 (0.0011) | 0.0011 (0.0009) |
| Observations | 118,081 | 118,081 | 118,081 | 118,081 | 118,081 |
| No of individuals | 6,299 | 6,299 | 6,299 | 6,299 | 6,299 |
| R-squared | 0.0037 | 0.0023 | 0.0013 | 0.0015 | 0.0016 |
| Mean $Y_{t=-1}$ | 0.00367 | 0.00080 | 0.00016 | 0.00032 | 0.00016 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20. All models contain year and age fixed effects.

Table A4a: Difference-in-Differences Event Study Models, Males

| VARIABLES | Father dies | | Mother dies | |
|-----------------------|-----------------------|----------------------|---------------------|-----------------------|
| | Treatment (1) | Control (2) | Treatment (3) | Control (4) |
| 8 years before death | 0.0003 (0.0008) | 0.0000 (0.0002) | -0.0006 (0.0015) | -0.0004 (0.0002) |
| 7 years before death | 0.0008 (0.0008) | -0.0001 (0.0002) | 0.0017 (0.0015) | -0.0004** (0.0002) |
| 6 years before death | 0.0008 (0.0007) | -0.0001 (0.0002) | -0.0006 (0.0014) | -0.0002 (0.0002) |
| 5 years before death | 0.0009 (0.0007) | 0.0001 (0.0002) | 0.0002 (0.0013) | -0.0003 (0.0002) |
| 4 years before death | 0.0006 (0.0007) | 0.0001 (0.0002) | -0.0003 (0.0012) | -0.0002 (0.0002) |
| 3 years before death | 0.0003 (0.0006) | 0.0001 (0.0002) | 0.0018 (0.0013) | -0.0001 (0.0002) |
| 2 years before death | 0.0006 (0.0006) | 0.0003* (0.0002) | 0.0006 (0.0011) | -0.0003* (0.0002) |
| Year of death | 0.0044*** (0.0008) | 0.0001 (0.0002) | 0.0028* (0.0015) | -0.0001 (0.0002) |
| 1 year after death | 0.0039*** (0.0009) | 0.0002 (0.0002) | 0.0020 (0.0015) | -0.0002 (0.0002) |
| 2 years after death | 0.0029*** (0.0009) | 0.0004* (0.0002) | 0.0001 (0.0015) | -0.0001 (0.0002) |
| 3 years after death | 0.0029*** (0.0010) | 0.0002 (0.0002) | 0.0002 (0.0016) | 0.0001 (0.0002) |
| 4 years after death | 0.0014 (0.0010) | 0.0004* (0.0003) | 0.0003 (0.0017) | 0.0000 (0.0003) |
| 5 years after death | 0.0021** (0.0010) | 0.0003 (0.0003) | -0.0008 (0.0018) | -0.0000 (0.0003) |
| 6 years after death | 0.0025** (0.0011) | 0.0003 (0.0003) | 0.0012 (0.0021) | 0.0001 (0.0003) |
| 7 years after death | 0.0023* (0.0012) | 0.0008** (0.0003) | 0.0006 (0.0022) | -0.0005 (0.0003) |
| 8 years after death | 0.0017 (0.0014) | 0.0003 (0.0003) | 0.0030 (0.0024) | 0.0003 (0.0004) |
| 9 years after death | 0.0025* (0.0015) | 0.0004 (0.0004) | 0.0012 (0.0027) | -0.0002 (0.0004) |
| 10 years after death | 0.0035** (0.0017) | 0.0005 (0.0004) | 0.0004 (0.0030) | -0.0004 (0.0004) |
| Observations | 343,600 | 2,614,608 | 125,187 | 2,769,002 |
| Number of individuals | 18,272 | 138,094 | 6,676 | 146,318 |
| R-squared | 0.0038 | 0.0022 | 0.0044 | 0.0022 |
| Mean $Y_{t=1}$ | 0.0045 | 0.0027 | 0.0062 | 0.0033 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. All models contain year and age fixed effects.

Table A4b: Difference-in-Differences Event Study Models, Females

| VARIABLES | Father dies | | Mother dies | |
|-----------------------|---------------------|-----------------------|-----------------------|---------------------|
| | Treatment (1) | Control (2) | Treatment (3) | Control (4) |
| 8 years before death | 0.0004 (0.0006) | 0.0002 (0.0002) | -0.0003 (0.0012) | 0.0000 (0.0002) |
| 7 years before death | 0.0006 (0.0006) | 0.0002 (0.0002) | 0.0010 (0.0012) | -0.0001 (0.0002) |
| 6 years before death | 0.0007 (0.0006) | 0.0001 (0.0002) | -0.0002 (0.0011) | -0.0000 (0.0002) |
| 5 years before death | 0.0003 (0.0006) | 0.0002 (0.0002) | 0.0003 (0.0011) | -0.0001 (0.0002) |
| 4 years before death | 0.0003 (0.0006) | 0.0001 (0.0002) | 0.0005 (0.0011) | -0.0002 (0.0002) |
| 3 years before death | -0.0003 (0.0006) | 0.0001 (0.0002) | 0.0013 (0.0011) | -0.0003 (0.0002) |
| 2 years before death | 0.0005 (0.0006) | 0.0002 (0.0002) | 0.0013 (0.0010) | -0.0002 (0.0002) |
| Year of death | 0.0004 (0.0006) | 0.0003 (0.0002) | 0.0039*** (0.0011) | -0.0002 (0.0002) |
| 1 year after death | 0.0004 (0.0007) | -0.0000 (0.0002) | 0.0033*** (0.0013) | 0.0000 (0.0002) |
| 2 years after death | 0.0014* (0.0008) | 0.0004* (0.0002) | 0.0028** (0.0014) | -0.0002 (0.0002) |
| 3 years after death | 0.0012 (0.0008) | 0.0002 (0.0002) | 0.0018 (0.0014) | 0.0001 (0.0002) |
| 4 years after death | 0.0012 (0.0009) | 0.0004* (0.0002) | 0.0006 (0.0015) | 0.0003 (0.0002) |
| 5 years after death | 0.0007 (0.0010) | 0.0009*** (0.0003) | 0.0008 (0.0016) | 0.0002 (0.0003) |
| 6 years after death | 0.0006 (0.0010) | 0.0006** (0.0003) | 0.0009 (0.0018) | 0.0003 (0.0003) |
| 7 years after death | 0.0005 (0.0011) | 0.0004 (0.0003) | 0.0004 (0.0019) | 0.0003 (0.0003) |
| 8 years after death | 0.0002 (0.0012) | 0.0008** (0.0003) | 0.0009 (0.0021) | 0.0004 (0.0003) |
| 9 years after death | 0.0008 (0.0012) | 0.0003 (0.0004) | 0.0008 (0.0023) | 0.0002 (0.0004) |
| 10 years after death | 0.0002 (0.0014) | 0.0002 (0.0004) | -0.0002 (0.0025) | 0.0003 (0.0004) |
| Observations | 334,878 | 2,475,950 | 118,081 | 2,636,944 |
| Number of individuals | 17,798 | 130,887 | 6,299 | 139,468 |
| R-squared | 0.0028 | 0.0015 | 0.0037 | 0.0014 |
| Mean $Y_{t=1}$ | 0.0039 | 0.0021 | 0.0037 | 0.0026 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. All models contain year and age fixed effects.

Table A5a: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death, Parental Death at Ages 10-15

| VARIABLES | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|-----------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | 0.0031* (0.0019) | 0.0032 (0.0033) | 0.0015 (0.0011) | -0.0013 (0.0028) |
| 7 years before death | 0.0031* (0.0017) | 0.0039 (0.0030) | 0.0015 (0.0010) | 0.0007 (0.0027) |
| 6 years before death | 0.0033** (0.0015) | 0.0007 (0.0026) | 0.0012 (0.0010) | -0.0015 (0.0023) |
| 5 years before death | 0.0015 (0.0013) | 0.0022 (0.0024) | 0.0004 (0.0009) | -0.0010 (0.0019) |
| 4 years before death | 0.0013 (0.0012) | 0.0012 (0.0022) | -0.0005 (0.0008) | 0.0009 (0.0018) |
| 3 years before death | 0.0012 (0.0010) | 0.0030 (0.0022) | -0.0005 (0.0007) | 0.0024 (0.0016) |
| 2 years before death | 0.0012 (0.0009) | -0.0013 (0.0016) | -0.0000 (0.0007) | 0.0017 (0.0012) |
| Year of death | 0.0040*** (0.0010) | -0.0003 (0.0019) | 0.0004 (0.0007) | 0.0017 (0.0011) |
| 1 year after death | 0.0047*** (0.0011) | 0.0004 (0.0022) | -0.0000 (0.0008) | 0.0032** (0.0016) |
| 2 years after death | 0.0035*** (0.0012) | -0.0045** (0.0020) | 0.0010 (0.0009) | 0.0040** (0.0020) |
| 3 years after death | 0.0034** (0.0014) | -0.0030 (0.0023) | 0.0009 (0.0011) | 0.0023 (0.0019) |
| 4 years after death | 0.0010 (0.0015) | -0.0010 (0.0026) | 0.0010 (0.0013) | 0.0006 (0.0025) |
| 5 years after death | 0.0015 (0.0018) | -0.0049 (0.0030) | 0.0016 (0.0015) | 0.0005 (0.0029) |
| 6 years after death | 0.0018 (0.0020) | -0.0003 (0.0033) | 0.0011 (0.0016) | 0.0019 (0.0031) |
| 7 years after death | 0.0024 (0.0022) | -0.0012 (0.0036) | 0.0019 (0.0019) | -0.0014 (0.0036) |
| 8 years after death | 0.0019 (0.0025) | -0.0013 (0.0044) | 0.0013 (0.0021) | 0.0001 (0.0041) |
| 9 years after death | 0.0028 (0.0029) | -0.0010 (0.0052) | 0.0003 (0.0023) | -0.0022 (0.0046) |
| 10 years after death | 0.0027 (0.0032) | -0.0015 (0.0054) | -0.0003 (0.0025) | -0.0078 (0.0050) |
| Observations | 155,805 | 53,561 | 152,659 | 51,734 |
| Number of individuals | 8,271 | 2,854 | 8,097 | 2,757 |
| R-squared | 0.0040 | 0.0046 | 0.0032 | 0.0052 |
| Mean Y_{t-1} | 0.00231 | 0.00565 | 0.00211 | 0.00146 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–15. All models contain year and age fixed effects.

Table A5b: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death, Parental Death at Ages 16-20

| VARIABLES | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|-----------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | 0.0022 (0.0016) | -0.0013 (0.0032) | -0.0010 (0.0015) | 0.0026 (0.0029) |
| 7 years before death | 0.0019 (0.0015) | 0.0015 (0.0029) | -0.0006 (0.0014) | 0.0029 (0.0029) |
| 6 years before death | 0.0013 (0.0013) | -0.0008 (0.0026) | -0.0003 (0.0014) | 0.0026 (0.0027) |
| 5 years before death | 0.0025** (0.0013) | -0.0013 (0.0021) | -0.0006 (0.0014) | 0.0026 (0.0026) |
| 4 years before death | 0.0015 (0.0011) | -0.0016 (0.0020) | 0.0001 (0.0013) | 0.0011 (0.0024) |
| 3 years before death | 0.0006 (0.0010) | 0.0010 (0.0018) | -0.0008 (0.0011) | 0.0010 (0.0022) |
| 2 years before death | 0.0006 (0.0009) | 0.0021 (0.0016) | 0.0004 (0.0010) | 0.0018 (0.0018) |
| Year of death | 0.0038*** (0.0012) | 0.0052** (0.0022) | 0.0012 (0.0010) | 0.0046** (0.0018) |
| 1 year after death | 0.0014 (0.0015) | 0.0032 (0.0025) | 0.0023* (0.0014) | 0.0019 (0.0019) |
| 2 years after death | -0.0001 (0.0018) | 0.0032 (0.0031) | 0.0040** (0.0016) | 0.0009 (0.0024) |
| 3 years after death | -0.0005 (0.0023) | 0.0020 (0.0035) | 0.0047** (0.0020) | -0.0003 (0.0029) |
| 4 years after death | -0.0016 (0.0025) | 0.0009 (0.0039) | 0.0053** (0.0024) | -0.0012 (0.0035) |
| 5 years after death | -0.0014 (0.0027) | 0.0022 (0.0044) | 0.0045 (0.0028) | -0.0000 (0.0042) |
| 6 years after death | -0.0013 (0.0030) | 0.0033 (0.0052) | 0.0056* (0.0031) | -0.0010 (0.0046) |
| 7 years after death | -0.0027 (0.0033) | 0.0036 (0.0057) | 0.0056* (0.0034) | -0.0002 (0.0051) |
| 8 years after death | -0.0045 (0.0037) | 0.0083 (0.0064) | 0.0062* (0.0037) | -0.0007 (0.0058) |
| 9 years after death | -0.0043 (0.0041) | 0.0061 (0.0071) | 0.0084** (0.0039) | 0.0001 (0.0066) |
| 10 years after death | -0.0042 (0.0046) | 0.0051 (0.0080) | 0.0088** (0.0043) | 0.0013 (0.0075) |
| Observations | 187,795 | 71,626 | 182,219 | 66,347 |
| Number of individuals | 10,001 | 3,822 | 9,701 | 3,542 |
| R-squared | 0.0040 | 0.0046 | 0.0026 | 0.0035 |
| Mean Y_{t-1} | 0.00622 | 0.00657 | 0.00539 | 0.00539 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 16–20. All models contain year and age fixed effects.

Table A6a: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death, Parent without Post-Compulsory Education

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|-----------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | 0.0019 (0.0012) | -0.0003 (0.0025) | 0.0003 (0.0009) | -0.0028 (0.0021) |
| 7 years before death | 0.0023* (0.0012) | 0.0018 (0.0024) | 0.0012 (0.0009) | 0.0003 (0.0022) |
| 6 years before death | 0.0023** (0.0011) | -0.0021 (0.0021) | 0.0007 (0.0009) | -0.0020 (0.0020) |
| 5 years before death | 0.0012 (0.0011) | 0.0000 (0.0020) | -0.0002 (0.0009) | -0.0011 (0.0019) |
| 4 years before death | 0.0004 (0.0010) | -0.0010 (0.0020) | 0.0009 (0.0010) | 0.0001 (0.0019) |
| 3 years before death | 0.0011 (0.0010) | 0.0005 (0.0020) | 0.0004 (0.0009) | 0.0005 (0.0019) |
| 2 years before death | 0.0007 (0.0009) | 0.0000 (0.0019) | 0.0003 (0.0009) | 0.0009 (0.0018) |
| Year of death | 0.0043*** (0.0012) | 0.0042* (0.0024) | 0.0006 (0.0009) | 0.0039** (0.0018) |
| 1 year after death | 0.0045*** (0.0013) | 0.0024 (0.0024) | -0.0003 (0.0010) | 0.0025 (0.0019) |
| 2 years after death | 0.0050*** (0.0014) | 0.0018 (0.0023) | 0.0014 (0.0011) | 0.0043* (0.0023) |
| 3 years after death | 0.0027* (0.0014) | 0.0021 (0.0025) | -0.0004 (0.0011) | 0.0017 (0.0022) |
| 4 years after death | 0.0026* (0.0015) | 0.0018 (0.0029) | 0.0006 (0.0012) | 0.0006 (0.0024) |
| 5 years after death | 0.0019 (0.0015) | 0.0006 (0.0029) | 0.0004 (0.0014) | -0.0003 (0.0027) |
| 6 years after death | 0.0025 (0.0017) | 0.0043 (0.0034) | 0.0004 (0.0015) | -0.0009 (0.0029) |
| 7 years after death | 0.0012 (0.0018) | 0.0033 (0.0035) | -0.0007 (0.0016) | -0.0020 (0.0030) |
| 8 years after death | 0.0006 (0.0021) | 0.0032 (0.0038) | -0.0013 (0.0017) | -0.0020 (0.0033) |
| 9 years after death | 0.0020 (0.0023) | 0.0038 (0.0043) | -0.0013 (0.0018) | 0.0011 (0.0039) |
| 10 years after death | 0.0029 (0.0026) | -0.0006 (0.0046) | -0.0009 (0.0019) | -0.0021 (0.0043) |
| Observations | 176,381 | 59,598 | 170,822 | 55,997 |
| Number of individuals | 9,382 | 3,178 | 9,077 | 2,981 |
| R-squared | 0.0039 | 0.0061 | 0.0032 | 0.0046 |
| Mean $Y_{t=1}$ | 0.00503 | 0.00729 | 0.00443 | 0.00540 |

Notes: Standard errors clustered at the individual level in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–15, only parents without post-compulsory education. All models contain year and age fixed effects.

Table A6b: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death, Parent with Post-Compulsory Education

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|-----------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | -0.0012 (0.0010) | -0.0010 (0.0018) | 0.0004 (0.0008) | 0.0019 (0.0012) |
| 7 years before death | -0.0007 (0.0010) | 0.0016 (0.0019) | 0.0000 (0.0007) | 0.0015 (0.0011) |
| 6 years before death | -0.0007 (0.0009) | 0.0008 (0.0018) | 0.0008 (0.0008) | 0.0014 (0.0010) |
| 5 years before death | 0.0005 (0.0010) | 0.0004 (0.0016) | 0.0009 (0.0009) | 0.0016 (0.0011) |
| 4 years before death | 0.0008 (0.0009) | 0.0003 (0.0014) | -0.0004 (0.0008) | 0.0008 (0.0011) |
| 3 years before death | -0.0004 (0.0008) | 0.0030* (0.0018) | -0.0009 (0.0007) | 0.0019 (0.0012) |
| 2 years before death | 0.0005 (0.0008) | 0.0011 (0.0013) | 0.0008 (0.0008) | 0.0016 (0.0010) |
| Year of death | 0.0045*** (0.0011) | 0.0016 (0.0017) | 0.0001 (0.0008) | 0.0039*** (0.0013) |
| 1 year after death | 0.0032*** (0.0011) | 0.0016 (0.0019) | 0.0012 (0.0010) | 0.0039** (0.0017) |
| 2 years after death | 0.0007 (0.0011) | -0.0013 (0.0020) | 0.0014 (0.0011) | 0.0015 (0.0016) |
| 3 years after death | 0.0029** (0.0012) | -0.0014 (0.0019) | 0.0028** (0.0012) | 0.0019 (0.0017) |
| 4 years after death | 0.0001 (0.0013) | -0.0011 (0.0020) | 0.0017 (0.0013) | 0.0007 (0.0018) |
| 5 years after death | 0.0021 (0.0014) | -0.0021 (0.0022) | 0.0009 (0.0013) | 0.0018 (0.0018) |
| 6 years after death | 0.0023 (0.0015) | -0.0017 (0.0026) | 0.0008 (0.0014) | 0.0027 (0.0021) |
| 7 years after death | 0.0034** (0.0017) | -0.0020 (0.0028) | 0.0017 (0.0016) | 0.0027 (0.0023) |
| 8 years after death | 0.0026 (0.0018) | 0.0027 (0.0031) | 0.0017 (0.0017) | 0.0038 (0.0026) |
| 9 years after death | 0.0030 (0.0020) | -0.0013 (0.0034) | 0.0029* (0.0018) | 0.0007 (0.0026) |
| 10 years after death | 0.0039* (0.0022) | 0.0008 (0.0039) | 0.0013 (0.0020) | 0.0016 (0.0028) |
| Observations | 167,219 | 65,589 | 164,056 | 62,084 |
| Number of individuals | 8,890 | 3,498 | 8,721 | 3,318 |
| R-squared | 0.0042 | 0.0047 | 0.0030 | 0.0044 |
| Mean $Y_{t=1}$ | 0.00384 | 0.00517 | 0.00334 | 0.00212 |


Notes: Standard errors clustered at the individual level in parentheses. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20, only parents with post-compulsory education. All models contain year and age fixed effects.

Table A7: Event Study Coefficients for Hospitalization, By Gender of Individual and by Parental Death, Family Fixed Effects Regressions

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|-----------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 8 years before death | -0.0006 (0.0023) | 0.0029 (0.0038) | -0.0028 (0.0021) | -0.0024 (0.0019) |
| 7 years before death | 0.0000 (0.0020) | 0.0047 (0.0034) | -0.0021 (0.0018) | -0.0009 (0.0018) |
| 6 years before death | 0.0003 (0.0017) | 0.0020 (0.0029) | -0.0016 (0.0015) | -0.0018 (0.0015) |
| 5 years before death | 0.0004 (0.0015) | 0.0023 (0.0024) | -0.0015 (0.0013) | -0.0010 (0.0014) |
| 4 years before death | 0.0003 (0.0012) | 0.0013 (0.0020) | -0.0011 (0.0011) | -0.0005 (0.0013) |
| 3 years before death | 0.0001 (0.0009) | 0.0028* (0.0017) | -0.0012 (0.0008) | 0.0006 (0.0012) |
| 2 years before death | 0.0005 (0.0007) | 0.0011 (0.0012) | 0.0001 (0.0007) | 0.0009 (0.0010) |
| Year of death | 0.0046*** (0.0009) | 0.0023 (0.0016) | 0.0008 (0.0007) | 0.0042*** (0.0012) |
| 1 year after death | 0.0042*** (0.0011) | 0.0009 (0.0018) | 0.0013 (0.0009) | 0.0039*** (0.0014) |
| 2 years after death | 0.0033*** (0.0013) | -0.0014 (0.0023) | 0.0028** (0.0011) | 0.0037** (0.0015) |
| 3 years after death | 0.0034** (0.0015) | -0.0018 (0.0027) | 0.0030** (0.0013) | 0.0030* (0.0016) |
| 4 years after death | 0.0021 (0.0018) | -0.0022 (0.0032) | 0.0034** (0.0016) | 0.0021 (0.0018) |
| 5 years after death | 0.0028 (0.0020) | -0.0039 (0.0037) | 0.0033* (0.0019) | 0.0028 (0.0020) |
| 6 years after death | 0.0034 (0.0024) | -0.0025 (0.0043) | 0.0038* (0.0021) | 0.0031 (0.0022) |
| 7 years after death | 0.0034 (0.0027) | -0.0038 (0.0049) | 0.0041* (0.0025) | 0.0030 (0.0025) |
| 8 years after death | 0.0028 (0.0030) | -0.0019 (0.0055) | 0.0042 (0.0027) | 0.0037 (0.0027) |
| 9 years after death | 0.0039 (0.0034) | -0.0044 (0.0061) | 0.0051* (0.0030) | 0.0038 (0.0030) |
| 10 years after death | 0.0050 (0.0036) | -0.0060 (0.0068) | 0.0049 (0.0032) | 0.0033 (0.0032) |
| Observations | 343,596 | 125,186 | 334,870 | 118,077 |
| Number of individuals | 18,268 | 6,675 | 17,790 | 6,295 |
| R-squared | 0.1303 | 0.1449 | 0.1413 | 0.1591 |
| Age fixed effects | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES |
| Parent fixed effects | YES | YES | YES | YES |
| Additional controls | NO | NO | NO | NO |
| Mean $Y_{t=1}$ | 0.00445 | 0.00618 | 0.00389 | 0.00367 |


Notes: Standard errors clustered at the individual level in parentheses. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$. Each column is from a separate linear probability model where the dependent variable is a dummy variable equal to one for being hospitalized for a mental health condition in that year. Sample is limited to individuals with a parental death at ages 10–20. Family fixed effects account for time-invariant effects related to the parent who has died at time 0.

Figure A1: Hospitalization Rate Before and After the Parental Death in the Treatment Groups (Raw Data)




Notes: Figure shows the annual hospitalization rates due to mental health disorders before and after parental death at time 0. Figures are given separately for male and female children and by gender of the deceased parent. Sample is limited to individuals born in 1971–1986 with a parental death at ages 10–20.

Figure A2b: Event Study Results by Cause of Hospitalization, Males




Notes: The figures plot the coefficient estimates from event study regressions together with 95% confidence intervals (standard errors clustered at the individual level). See Tables A3a and A3b for tabulation of estimation results and further details.

Figure A2b: Event Study Results by Cause of Hospitalization, Females



Notes: The figures plot the coefficient estimates from event study regressions together with 95% confidence intervals (standard errors clustered at the individual level). See Tables A3c and A3d for tabulation of estimation results and further details.


Figure A3: Event Study Results, Intentional Self-Harm, Males and Females




Notes: The figures plot the coefficient estimates from event study regressions together with 95% confidence intervals (standard errors clustered at the individual level). See Tables A3a–A3d for tabulation of estimation results and further details.

Figure A4: Event Study Results by Parental Education

Panel A: Without post-compulsory education



Panel B: With post-compulsory education



Notes: The figures plot the coefficient estimates from event study regressions, together with 95% confidence intervals (standard errors clustered at the individual level). The outcome is the hospitalization rate. Panels on the left show estimates for father’s death and those on the right show them for mother’s death. Parental education refers to the education level of the parent who has died at time 0.

APPENDIX B. PRE-TREND AND IMPUTATION ESTIMATION RESULTS

Pre-Trend Testing for Our Baseline Model

To investigate the validity of the parallel trend assumption of our baseline event study approach, we follow Borusyak et al. (2021) and estimate our model using the set of observations prior to treatment:

$$Y_{ist} = \sum_{j=-7}^{-1} \gamma_j \cdot I[t = j] + \sum_k \beta_k \cdot I[age_{is} = k] + \sum_y \tau_y \cdot I[s = y] + \varepsilon_{ist},$$

where $I[t = j]$ are indicator variables of being treated 1 to 7 years later, the comparison group consisting of those experiencing the parental death eight years later. After estimation of the model, the joint statistical significance of the γ_j 's is tested using F-test. Further, the individual pre-trend coefficients can reveal possible anticipation effects to parental death that would violate the identification assumption of our model. As explained by Borusyak et al. (2021) the key advantage of this test approach is that it separates the validation of the parallel trend assumption from the estimation, given the event study design. The results reported in Table B1 do not reveal evidence of significant pre-trends based on individual t-tests or a joint significance F-test. Table B2 reports the corresponding results for the control group used in our difference-in-differences design. Again, we do not find evidence of significant pre-trends.

Table B1: Pre-Trend Testing for the Baseline Event Study Approach – Treatment Group

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|--------------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 7 years before death | 0.0004 (0.0006) | 0.0022** (0.0010) | 0.0003 (0.0004) | 0.0012 (0.0007) |
| 6 years before death | 0.0004 (0.0006) | 0.0000 (0.0011) | 0.0003 (0.0005) | 0.0001 (0.0006) |
| 5 years before death | 0.0006 (0.0007) | 0.0009 (0.0012) | -0.0002 (0.0005) | 0.0007 (0.0008) |
| 4 years before death | 0.0003 (0.0007) | 0.0003 (0.0012) | -0.0002 (0.0005) | 0.0010 (0.0009) |
| 3 years before death | 0.0000 (0.0007) | 0.0026* (0.0015) | -0.0007 (0.0005) | 0.0016 (0.0011) |
| 2 years before death | -0.0000 (0.0007) | 0.0016 (0.0014) | 0.0002 (0.0006) | 0.0013 (0.0012) |
| 1 years before death | -0.0003 (0.0008) | 0.0022 (0.0015) | -0.0001 (0.0007) | 0.0005 (0.0012) |
| Observations | 145,711 | 53,130 | 141,907 | 50,140 |
| Number of individuals | 18,259 | 6,669 | 17,788 | 6,289 |
| R-squared | 0.0013 | 0.0014 | 0.0021 | 0.0021 |
| Age fixed effects | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES |
| Individual fixed effects | NO | NO | NO | NO |

| | | | | |
|----------------|-------|-------|-------|-------|
| F-test value | 0.319 | 1.547 | 0.740 | 0.691 |
| F-test p-value | 0.946 | 0.147 | 0.638 | 0.680 |

Notes: The outcome is the annual hospitalization rate due to mental health disorders. Standard errors clustered at the individual level are given in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Comparison group is the 8th year before parental death. Pre-trend coefficients that have been estimated using OLS on the pre-treatment observations only. F-test reports the joint significance of the pre-treatment coefficients (-7, ..., -1).

Table B2: Pre-Trend Testing for the Event Study Approach – Control Group

| | Males, Father (1) | Males, Mother (2) | Females, Father (3) | Females, Mother (4) |
|--------------------------|-------------------------|-------------------------|---------------------------|---------------------------|
| 7 years before death | -0.0001 (0.0002) | -0.0001 (0.0002) | -0.0001 (0.0001) | -0.0002 (0.0001) |
| 6 years before death | -0.0001 (0.0002) | 0.0002 (0.0002) | -0.0002 (0.0001) | -0.0001 (0.0001) |
| 5 years before death | 0.0001 (0.0002) | 0.0001 (0.0002) | 0.0000 (0.0001) | -0.0001 (0.0001) |
| 4 years before death | 0.0001 (0.0002) | 0.0001 (0.0002) | -0.0001 (0.0002) | -0.0002 (0.0001) |
| 3 years before death | 0.0000 (0.0002) | 0.0002 (0.0002) | -0.0001 (0.0002) | -0.0003* (0.0002) |
| 2 years before death | 0.0003 (0.0002) | -0.0000 (0.0002) | -0.0001 (0.0002) | -0.0002 (0.0002) |
| 1 years before death | 0.0001 (0.0002) | 0.0004* (0.0002) | -0.0002 (0.0002) | -0.0001 (0.0002) |
| Observations | 1,101,859 | 1,166,978 | 1,044,562 | 1,112,535 |
| Number of individuals | 138,021 | 146,217 | 130,820 | 139,398 |
| R-squared | 0.0006 | 0.0006 | 0.0007 | 0.0007 |
| Age fixed effects | YES | YES | YES | YES |
| Year fixed effects | YES | YES | YES | YES |
| Individual fixed effects | NO | NO | NO | NO |
| F-test value | 0.671 | 1.588 | 0.660 | 0.728 |
| F-test p-value | 0.697 | 0.134 | 0.707 | 0.649 |

Notes: Sample includes control individuals without a parental death by age 30. The outcome is the annual hospitalization rate due to mental health disorders. Standard errors clustered at the individual level are given in parentheses. *** p<0.01, ** p<0.05, * p<0.1. Comparison group is the 8th year before parental death. Pre-trend coefficients that have been estimated using OLS on the pre-treatment observations only. F-test reports the joint significance of the pre-treatment coefficients (-7, ..., -1).

Imputation Estimator Results

We examine the robustness of our baseline model to difference-in-differences design with staggered adoption of treatment (see Borusyak et al., 2021, and an application by von Bismarck-Osten et al., 2020). The design assumes that the true causal model for individual i in year t is:

$$Y_{it} = \alpha_i + \beta_t + \tau_{it} \cdot I[t \geq E_i] + \varepsilon_{it},$$

where α_i are the individual fixed effects, β_t are the age fixed effects, E_i is the individual's age when a parent dies (i.e., treatment), and $I[t \geq E_i]$ is the indicator for the post-treatment periods. Here, τ_{it} captures the heterogeneous treatment effects, i.e., the effects of the death of father (or mother) on mental health-related hospitalizations.


The treatment effects can be estimated using the imputation estimator proposed by Borusyak et al. (2021). The imputation estimator is based on the parallel trends assumption, but contrary to the fixed effects OLS regression with lags and leads of treatment, it produces estimates that are robust even in the presence of heterogeneous treatment effects. The imputation estimator with a particular horizon h (i.e., h years after treatment) leverages all difference-in-differences contrasts between individual i in period $E_i + h$ relative to all periods before treatment, $t < E_i$, and relative to other individuals who have not been treated by $E_i + h$.

To validate the assumptions of the imputation estimator, Figure B1 reports first the estimated pre-trend coefficients, together with their 95% confidence intervals (depicted in red). The pre-trend coefficients have been estimated using OLS with individual and age fixed effects, because the imputation estimator utilizes individual fixed effects, rather than year fixed effects. Thus, these pre-trend estimates differ slightly from those reported in Table B1.¹⁸ However, the conclusion remains intact: the pre-trend coefficients are small and statistically insignificant, which provides support for the parallel trend assumption (and for the use of pre-treatment periods as the comparison group in the imputation estimation).

In Figure B1, treatment effects of parental death have been estimated using the imputation estimator (depicted in blue). In all cases, the estimated effects are similar to our baseline results reported in Figure 1. The pre-trend and imputation estimation results for the control group are shown in Figure B2. They are similar to those reported in Figure 2.


¹⁸ When individual fixed effects are used, we need to drop two time effects from the model for identification.

Figure B1: Pre-Trend Coefficients and Treatment Estimates – Treatment Group



Notes: The outcome is the annual hospitalization rate for mental health disorders. The OLS estimates depicted in red report pre-trend coefficients estimated using pre-treatment observations only. This model includes individual and age fixed effects and assumes zero effects 6–8 years prior to parental death. The treatment effects for years 0–8 depicted in blue have been estimated using the imputation estimator of Borusyak et al. (2021) that includes individual and age fixed effects. 95% confidence intervals are reported, based on standard errors clustered at the individual level.

Figure B2: Pre-Trend Coefficients and Treatment Estimates – Control Group



Notes: Sample includes control individuals without a parental death by age 30. The outcome is the annual hospitalization rate for mental health disorders. The OLS estimates depicted in red report pre-trend coefficients estimated using pre-treatment observations only. This model includes individual and age fixed effects and assumes zero effects 6–8 years prior to parental death. The treatment effects for years 0–8 depicted in blue have been estimated using the imputation estimator of Borusyak et al. (2021) that includes individual and age fixed effects. 95% confidence intervals are reported, based on standard errors clustered at the individual level.

APPENDIX C. UNINFORMATIVE VS. INFORMATIVE PARENTAL DEATHS

To identify the explicitly exogenous causes of death, we use the approach introduced by Espinosa and Evans (2008) and adopted later in Gimenez et al. (2013). The basic idea of this approach is to classify the causes of death into two non-overlapping groups: i) deaths strongly correlated with measures of parental SES (i.e., informative causes of death, ICOD) and ii) deaths driven by likely random causes and not correlated with SES (i.e., uninformative causes of death, UCOD).

In our data, the causes of deaths are recorded using the International Classification of Diseases (ICD-8 in 1969–1986, ICD-9 in 1987–1995 and ICD-10 since 1996). To identify the deaths that are informative and uninformative, we first regroup the causes of deaths ICDs according to the Statistics Finland’s classification of deaths into 54 subgroups (COD). We then use ordinary least squares estimation (OLS) to categorize each COD group according to its degree of correlation with family SES (income and education).

Following Gimenez et al. (2013), we estimate linear probability models for each of the 54 COD groups:

$$COD_{irt}^d = \alpha^d + \sum_{k=2}^4 \beta_k^d \cdot I[INC(k)_{irt}^d = 1] + \sum_{k=2}^4 \gamma_k^d \cdot I[EDU(k)_{irt}^d = 1] + \sum_{k=2}^4 \eta_k^d \cdot I[EDUS(k)_{irt}^d = 1] + \delta_r^d + \tau_t^d + X_{irt}^d \theta^d + \varepsilon_{irt}^d$$

where COD_{irt}^d is 1 for parent i who resided in region r and died from cause of death d in year t (and 0 otherwise). For example, variable COD_{irt}^6 equals 1 if a parent i died from a malignant neoplasm of stomach during a 25-year follow-up period after the child’s birth, and 0 if the cause of death is different. $I[INC(k)_{irt}^d = 1]$ equals one if the family income of parent i is in the k^{th} income quartile; $I[EDU(k)_{irt}^d = 1]$ and $I[EDUS(k)_{irt}^d = 1]$ represent four indicator variables for the education level k attained by the deceased parent i and attained by his/her spouse, respectively. δ_r^d and τ_t^d represent region and year-of-death fixed effects, respectively. X_{irt}^d contains four dummies of age (quartiles) at the time of death. Models are estimated separately for fathers and mothers.

For each regression, we conducted four F-tests: i) whether the coefficient estimates for the income level, ii) the education indicators for the deceased, iii) the education indicators for the spouse of the deceased, and iv) all three sets, income, own education, and spousal education indicators, respectively, are jointly zero. If we can reject any of the four null hypotheses at the five percent confidence level, then the cause of death COD^d is considered to be informative (ICOD); otherwise, the COD^d is considered to be uninformative (UCOD).

Table C1 summarizes the results of the regressions for top five ICODs and UCODs for fathers and mothers. Overall, approximately 14.7 percent of the father’s deaths were classified as UCODs. The corresponding figure for the mothers is 8.9 percent. To evaluate the robustness of our baseline findings, we estimate separate effects for uninformative causes of death (UCOD) and informative causes of death (ICOD) using event study models and cross-sectional models. These results are illustrated in Figures 3 and C1.


Table C1: Leading Informative and Uninformative Causes of Death

| <i>Informative Causes of Death (ICOD)</i> | Mortality rate | P-value on F-test that the estimates are jointly zero | | | |
|--|----------------|---|------------------|---------------|-------|
| | | Father education | Mother education | Family income | All |
| <i>Death of a Father</i> | | | | | |
| Ischaemic heart diseases | 15.367 | 0.000 | 0.000 | 0.001 | 0.000 |
| Suicides | 10.917 | 0.000 | 0.000 | 0.000 | 0.000 |
| Alcohol-related diseases and accidental poisoning by alcohol | 10.141 | 0.000 | 0.069 | 0.342 | 0.000 |
| Malignant neoplasm of larynx, trachea, bronchus and lung | 3.231 | 0.000 | 0.000 | 0.091 | 0.000 |
| Other malignant neoplasms | 2.900 | 0.000 | 0.000 | 0.001 | 0.000 |
| <i>Death of a Mother</i> | | | | | |
| Malignant neoplasm of breast | 4.242 | 0.004 | 0.000 | 0.000 | 0.000 |
| Suicides | 2.693 | 0.537 | 0.530 | 0.000 | 0.004 |
| Alcohol-related diseases and accidental poisoning by alcohol | 2.473 | 0.042 | 0.000 | 0.000 | 0.000 |
| Other malignant neoplasms | 2.303 | 0.059 | 0.000 | 0.584 | 0.000 |
| Cerebrovascular diseases | 2.115 | 0.063 | 0.002 | 0.001 | 0.000 |

| <i>Uninformative Causes of Death (UCOD)</i> | Mortality rate | P-value on F-test that the estimates are jointly zero | | | |
|---|----------------|---|------------------|---------------|-------|
| | | Father education | Mother education | Family income | All |
| <i>Death of a Father</i> | | | | | |
| Cerebrovascular diseases | 4.409 | 0.349 | 0.671 | 0.849 | 0.787 |
| Other heart diseases excl. rheumatic and alcohol-related | 2.502 | 0.136 | 0.708 | 0.640 | 0.511 |
| Accidental falls | 2.134 | 0.113 | 0.227 | 0.777 | 0.303 |
| Diabetes mellitus | 0.921 | 0.844 | 0.132 | 0.428 | 0.364 |
| Event of undetermined intent | 0.753 | 0.533 | 0.471 | 0.291 | 0.444 |
| <i>Death of a Mother</i> | | | | | |
| Land traffic accidents | 1.021 | 0.556 | 0.397 | 0.114 | 0.198 |
| Malignant melanoma of skin | 0.296 | 0.378 | 0.317 | 0.051 | 0.075 |
| Malignant neoplasm of rectum, anus and anal canal | 0.258 | 0.375 | 0.309 | 0.707 | 0.553 |
| Malignant neoplasm of kidney | 0.232 | 0.133 | 0.480 | 0.788 | 0.497 |
| Primary malignant neoplasm of liver and intrahepatic bile ducts | 0.224 | 0.415 | 0.194 | 0.551 | 0.585 |

Notes: The top five CODs’ mortality rates are given. Mortality rate is measured as number of deaths per 1,000 individuals.

Figure C1: Cross-Sectional Linear Probability Model for Hospitalization at Ages 26–30, Uninformative Causes of Death (Males and Females)



Notes: Figure reports coefficient estimates together with 95% confidence intervals (based on robust standard errors). Comparison group is the individuals without parental death by age 30. Coefficients are for uninformative causes of deaths. Model also include unreported coefficients for informative causes of deaths.

APPENDIX D. TABLES AND FIGURES: CROSS-SECTIONAL ANALYSIS

To measure sickness absenteeism, we use total data on sickness absence spells and days over the period 1995–2016. The comprehensive register-based data are from the Social Insurance Institution of Finland (Kela). Information is derived from the database used to pay the sickness allowances to the affected persons. Sickness absences are diagnosed by a physician (i.e., general physicians, occupational physicians, and psychiatrists). Before receiving the sickness allowance from Kela, the person must complete a nine-day waiting period. The incapacity for work must be certified by a doctor, and the employer is obliged to notify Kela of the sickness leave. The employee is entitled to the normal, full salary during the nine-day waiting period (for a detailed description of the Finnish sickness insurance system, see Böckerman et al., 2018). Thus, due to the institutional character of the system, the data recorded by Kela contain sickness absence spells lasting longer than nine days. The medical reason for sickness leave is comprehensively recorded for the period 2004–2016, allowing us to identify mental health-related sickness leaves.

To capture mild mental health-related disorders that do not lead to hospitalization, we utilize data from the Social Insurance Institution of Finland containing filled mental health-related medications dispensed at Finnish pharmacies over 1995–2016. These medications are listed in the World Health Organization (WHO)'s anatomical therapeutic chemical (ATC) classification system as codes beginning with 'N05', 'N06A', 'N06B' or 'N06C'. As anti-depressants are the first choice of treatment in moderate and severe depression cases in Finland based on the clinical guidelines, including the use of mental health-related medications in the set of outcomes also captures individuals who are not treated in hospital or outpatient clinics and who did not need sick absence leave due to their depression or other mental disorder. The data record the universe of individual-level prescriptions reimbursed under comprehensive national health insurance scheme. All permanent residents of Finland are automatically covered under the national health insurance. The use of comprehensive

register-based data allows us to follow patients over time, even in cases where patients switch physicians or employers. We create an indicator of having at least one prescription for mental health-related disorders per year.

We describe the cross-sectional results for non-economic and economic consequences of parental death more fully in the following. First, we present results for the baseline dependent variable that equals one for individuals who were hospitalized for a mental health-related condition between the ages of 26 and 30. These results are reported in columns 2 and 4 in Table D2 and Figure D1; see also Figures 4a and 4b. The most striking result is that the parental death variables are often large and statistically significant, regardless of the gender of the child or the parent. In other words, the short-run pattern, from the event studies, where coefficients are the largest for the death of the parent of the same gender, is not present in the long-run, cumulative analysis using cross-sectional data. The coefficients are usually larger for males than females. When comparing the effects at different ages of parental deaths, we observe more similarities than differences either for males (left panel) or females (right panel). The results for uninformative causes of deaths are similar (see Figure C1).

To obtain a more comprehensive picture of the health effects, we proceed to analyze the relationship between parental death and several additional cross-sectional outcomes that can be measured only in adulthood. The results, shown in Figures 4a and 4b, are based on the same cross-sectional model (equation 2). An event study approach, based on pre- and post-death outcomes, is not appropriate for outcomes that are not relevant for children, such as work-related absences.

The overall hospitalization measure used in our main estimation results does not capture mild mental health-related problems not requiring hospitalization in specialized public health care. For this reason, we examine separately the relationship between parental death and the use of mental health-related medications at ages 29–30. We find a significant increase in the use of mental health-related medications after parental death, for both males

and females. The effect among both genders is interesting for two reasons. First, the use of mental health-related medications is more common among females in the population. Second, the increase in the use of mental health-related medications does not eliminate the appearance of serious mental health-related problems captured by the hospitalization measure.

The results also show a substantial increase in mental health-related deaths at ages 26–30 for males due to maternal death. Further, we observe an increase in sickness absence days at ages 26–30 after parental death prevailing among both the genders. The quantitative size of the effect is larger for females. There is also an increase in sickness absence days due to mental health-related reasons for both genders at age 30.

Finally, using register-based data, we study whether broad economic outcomes in adulthood are negatively affected by parental death. We find that there are substantial negative effects on education (measured by years of schooling), annual earnings at ages 26–30 and employment, for both females and males (Figure D2). These effects are not notably different due to paternal or maternal death. The effects on two key labor market outcomes (i.e., earnings and employment) are larger for males. The larger effects for males are most likely related to their stronger labor market attachment and higher average earnings. These significant negative effects on income level and socio-economic status in adulthood could partially lead to poor mental health.

Table D1: Mean Values by Parental Death, Cross-Sectional Analysis

| Parental death | No parental death by age 30 | Father's death by age 30 | Mother's death by age 30 |
|---|-----------------------------|--------------------------|--------------------------|
| <i>Dependent variables</i> | | | |
| Hospitalization spell due to mental health disorders (0/1; ages 26–30) | 0.0202 | 0.0327 | 0.0368 |
| Hospitalization spell due to depression (0/1; ages 26–30) | 0.00539 | 0.0081 | 0.0099 |
| Hospitalization spell due to anxiety (0/1; ages 26–30) | 0.00202 | 0.0031 | 0.0035 |
| Hospitalization spell due to substance-use disorder (0/1; ages 26–30) | 0.00553 | 0.0122 | 0.0134 |
| Hospitalization spell due to intentional self-harm (0/1; ages 26–30) | 0.00332 | 0.0065 | 0.0076 |
| Hospitalization day visit due to mental health disorders (0/1; ages 27–30) | 0.0483 | 0.0653 | 0.0735 |
| Mental health-related death (0/1; ages 26–30) | 0.000158 | 0.0049 | 0.0054 |
| Using mental health-related prescription drugs (0/1; ages 29–30) | 0.0943 | 0.116 | 0.125 |
| Reimbursed sickness absence days, days per annum (cont. variable; ages 26–30) | 2.116 | 2.700 | 2.812 |
| Sick leave due to mental health disorders (0/1; age 30) | 0.0141 | 0.0184 | 0.0224 |
| Years of schooling (by age 30) | 13.60 | 12.85 | 12.86 |
| Annual earnings (ages 26–30) | 22.68 | 19.81 | 19.94 |
| Employment rate (ages 26–30) | 0.761 | 0.697 | 0.693 |
| <i>Independent variables</i> | | | |
| Female (0/1) | 0.487 | 0.491 | 0.486 |
| Language Finnish (0/1) | 0.947 | 0.960 | 0.956 |
| Language Swedish (0/1) | 0.0516 | 0.0390 | 0.0428 |
| Other Language (0/1) | 0.0013 | 0.0010 | 0.0012 |
| Lived with father at childhood (0/1) | 0.923 | 0.821 | 0.870 |
| Lived with mother at childhood (0/1) | 0.971 | 0.961 | 0.918 |
| Father has only basic education (0/1) | 0.364 | 0.516 | 0.452 |
| Father has upper secondary educ. (0/1) | 0.376 | 0.318 | 0.342 |
| Father has tertiary education (0/1) | 0.260 | 0.166 | 0.206 |
| Mother has only basic education (0/1) | 0.368 | 0.473 | 0.491 |
| Mother has upper secondary educ. (0/1) | 0.391 | 0.358 | 0.328 |
| Mother has tertiary education (0/1) | 0.241 | 0.169 | 0.181 |
| Father completed high school (0/1) | 0.181 | 0.096 | 0.138 |
| Mother completed high school (0/1) | 0.257 | 0.177 | 0.174 |
| Father's taxable income percentile (0–1) | 0.592 | 0.556 | 0.583 |
| Mother's taxable income percentile (0–1) | 0.572 | 0.561 | 0.551 |
| Father has been hospitalized due to mental health disorder (0/1) | 0.0329 | 0.133 | 0.074 |
| Father has been hospitalized due to WAAC (0/1) | 0.0140 | 0.0913 | 0.045 |
| Mother has been hospitalized due to mental health disorder (0/1) | 0.0247 | 0.0491 | 0.114 |
| Mother has been hospitalized due to WAAC (0/1) | 0.00391 | 0.0135 | 0.0413 |
| Mother's age at child birth (0/1) | 27.12 | 28.69 | 29.23 |

| | | | |
|--------------------------------------|---------|---------|--------|
| No. of siblings w/ same mother (0/1) | 2.120 | 2.007 | 2.058 |
| Father missing (0/1) | 0.0166 | 0 | 0.0553 |
| Mother missing (0/1) | 0.00166 | 0.0353 | 0 |
| <hr/> | | | |
| Number of individuals | 818,923 | 112,280 | 41,840 |

Notes: Data also include indicator variables for birth year (1971–86), birth region (19 regions) and parents' occupation (9 categories for father and mother). Parents' income has been measured in 1970 & 1975 if the birth year is 1971–75, in 1975 & 1980 if it is 1976–81, and 1980 & 1985 if it is 1981–86. WAAC = Wholly alcohol-attributable conditions. Parents' hospitalization outcomes measured by child's age 9. Paternal (maternal) figures exclude individuals without missing information on the father (mother).


Table D2: Cross-Sectional Linear Probability Model, Hospitalization at Ages 26–30

| | Males (1) | Males (2) | Females (3) | Females (4) |
|----------------------------------|-----------------------|------------------------|-----------------------|------------------------|
| Age 0–6 at paternal death | 0.0196*** (0.0031) | 0.0071** (0.0032) | 0.0122*** (0.0027) | 0.0044 (0.0027) |
| Age 7–10 at paternal death | 0.0144*** (0.0031) | 0.0053* (0.0031) | 0.0140*** (0.0028) | 0.0086*** (0.0028) |
| Age 11–15 at paternal death | 0.0147*** (0.0023) | 0.0076*** (0.0023) | 0.0071*** (0.0019) | 0.0027 (0.0019) |
| Age 16–20 at paternal death | 0.0136*** (0.0019) | 0.0084*** (0.0019) | 0.0098*** (0.0017) | 0.0065*** (0.0017) |
| Age 21–25 at paternal death | 0.0128*** (0.0016) | 0.0086*** (0.0016) | 0.0075*** (0.0014) | 0.0051*** (0.0014) |
| Age 26–30 at paternal death | 0.0111*** (0.0014) | 0.0081*** (0.0014) | 0.0064*** (0.0012) | 0.0045*** (0.0012) |
| Age 0–6 at maternal death | 0.0175*** (0.0059) | 0.0094 (0.0059) | 0.0134** (0.0054) | 0.0052 (0.0054) |
| Age 7–10 at maternal death | 0.0198*** (0.0057) | 0.0111* (0.0057) | 0.0040 (0.0044) | -0.0020 (0.0044) |
| Age 11–15 at maternal death | 0.0229*** (0.0043) | 0.0167*** (0.0043) | 0.0071** (0.0034) | 0.0029 (0.0034) |
| Age 16–20 at maternal death | 0.0164*** (0.0032) | 0.0118*** (0.0032) | 0.0179*** (0.0032) | 0.0147*** (0.0032) |
| Age 21–25 at maternal death | 0.0182*** (0.0028) | 0.0143*** (0.0028) | 0.0085*** (0.0024) | 0.0058** (0.0023) |
| Age 26–30 at maternal death | 0.0171*** (0.0024) | 0.0139*** (0.0023) | 0.0072*** (0.0020) | 0.0047** (0.0019) |
| Swedish language | | -0.0084*** (0.0010) | | -0.0067*** (0.0009) |
| Other native language | | -0.0106* (0.0057) | | -0.0104** (0.0052) |
| Mother's age at child birth | | -0.0001** (0.0001) | | 0.0000 (0.0000) |
| Lived with father at childhood | | -0.0093*** (0.0011) | | -0.0069*** (0.0010) |
| Lived with mother at childhood | | 0.0029 (0.0018) | | -0.0035** (0.0017) |
| Father completed high school | | -0.0007 (0.0007) | | 0.0004 (0.0007) |
| Mother completed high school | | -0.0010 (0.0006) | | 0.0007 (0.0006) |
| Father has upper secondary educ. | | -0.0018*** (0.0005) | | -0.0002 (0.0005) |
| Father has tertiary education | | -0.0011 (0.0008) | | -0.0014** (0.0007) |
| Mother has upper secondary educ. | | -0.0039*** (0.0006) | | -0.0019*** (0.0005) |
| Mother has tertiary education | | -0.0033*** (0.0008) | | -0.0019*** (0.0007) |
| Father's income quartile 1/4 | | -0.0053*** (0.0010) | | -0.0042*** (0.0009) |
| Father's income quartile 2/4 | | -0.0082*** (0.0011) | | -0.0064*** (0.0010) |
| Father's income quartile 3/4 | | -0.0103*** (0.0011) | | -0.0066*** (0.0010) |
| Mother's income quartile 1/4 | | -0.0012 | | -0.0008 |

| | | | | |
|---|---------|------------|---------|------------|
| | | (0.0009) | | (0.0008) |
| Mother's income quartile 2/4 | | -0.0042*** | | -0.0043*** |
| | | (0.0009) | | (0.0008) |
| Mother's income quartile 3/4 | | -0.0055*** | | -0.0053*** |
| | | (0.0010) | | (0.0009) |
| Father has been hospitalized due to mental health disorder | | 0.0215*** | | 0.0215*** |
| | | (0.0019) | | (0.0019) |
| Father has been hospitalized due to WAAC | | 0.0067** | | 0.0067** |
| | | (0.0027) | | (0.0027) |
| Mother has been hospitalized due to mental health disorder | | 0.0349*** | | 0.0349*** |
| | | (0.0021) | | (0.0021) |
| Mother has been hospitalized due to WAAC | | 0.0037 | | 0.0037 |
| | | (0.0047) | | (0.0047) |
| No. of siblings w/ same mother | | 0.0007*** | | 0.0006*** |
| | | (0.0002) | | (0.0002) |
| Observations | 494,332 | 494,332 | 470,601 | 470,601 |
| R-squared | 0.0026 | 0.0087 | 0.0009 | 0.0050 |
| Controls | NO | YES | NO | YES |
| Mean hospitalization rate | 0.0244 | 0.0244 | 0.0196 | 0.0196 |


Notes: Outcome variable is a dummy for having at least one hospitalization spell due to mental health disorders at ages 26–30. Robust standard errors in parentheses. All regressions also contain controls for birth year, birth region, parents' occupation (9 categories for father and mother), unknown father and unknown mother.
*** p<0.01, ** p<0.05, * p<0.1 (for two-sided tests).

Figure D1a: Cross-Sectional Linear Probability Model, Alternate Health Outcomes, Males




Notes: Figure reports coefficient estimates together with 95% confidence intervals based on robust standard errors. Model specification is the same as in Appendix Table D2 (with controls). Comparison group is children born in 1971–86 without parental death by age 30.

Figure D1b: Cross-Sectional Linear Probability Model, Alternate Health Outcomes, Females



Notes: Figure reports coefficient estimates together with 95% confidence intervals based on robust standard errors. Model specification is the same as in Appendix Table D2 (with controls). Comparison group is children born in 1971–86 without parental death by age 30.

Figure D2: Cross-Sectional Linear Probability Model, Education and Labour Market Outcomes (Males and Females)



Notes: Figure reports coefficient estimates together with 95% confidence intervals based on robust standard errors. Model specification is the same as in Appendix Table D2 (with controls). Comparison group is children born in 1971–86 without parental death by age 30.