

DISCUSSION PAPER SERIES

IZA DP No. 10872

**Life After Lead:
Effects of Early Interventions for Children
Exposed to Lead**

Stephen B. Billings
Kevin T. Schnepel

JULY 2017

DISCUSSION PAPER SERIES

IZA DP No. 10872

Life After Lead: Effects of Early Interventions for Children Exposed to Lead

Stephen B. Billings

University of North Carolina at Charlotte

Kevin T. Schnepel

University of Sydney and IZA

JULY 2017

Any opinions expressed in this paper are those of the author(s) and not those of IZA. Research published in this series may include views on policy, but IZA takes no institutional policy positions. The IZA research network is committed to the IZA Guiding Principles of Research Integrity.

The IZA Institute of Labor Economics is an independent economic research institute that conducts research in labor economics and offers evidence-based policy advice on labor market issues. Supported by the Deutsche Post Foundation, IZA runs the world's largest network of economists, whose research aims to provide answers to the global labor market challenges of our time. Our key objective is to build bridges between academic research, policymakers and society.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ABSTRACT

Life After Lead: Effects of Early Interventions for Children Exposed to Lead*

Lead pollution is consistently linked to cognitive and behavioral impairments, yet little is known about the benefits of public health interventions for children exposed to lead. This paper estimates the long-term impacts of early-life interventions (e.g. lead remediation, nutritional assessment, medical evaluation, developmental surveillance, and public assistance referrals) recommended for lead-poisoned children. Using linked administrative data from Charlotte, NC, we compare outcomes for children who are similar across observable characteristics but differ in eligibility for intervention due to blood lead test results. We find that the negative outcomes previously associated with early-life exposure can largely be reversed by intervention.

JEL Classification: I12, I18, I21, J13, J24, K42, Q53, Q58

Keywords: early childhood intervention, early health shocks, lead exposure, human capital formation

Corresponding author:

Kevin T. Schnepel
H04 – Merewether
The University of Sydney
NSW 2006
Australia

E-mail: kevin.schnepel@sydney.edu.au

* We thank Anna Aizer, Alan Barreca, Prashant Bharadwaj, David Card, Deborah Cobb-Clark, Susan Dynarski, Nathaniel Hendren, Jason Lindo, John Lynch, Bhash Mazumder, Mike Mueller-Smith, Trevor O'Grady, Jessica Wolpaw Reyes, Heather Royer, Lisa Schulkind, and Barbara Wolfe for helpful comments/discussions as well as seminar participants at the 2014 Labour and Econometrics Workshop at the University of Otago, the 2015 IZA 7th Annual Meeting on the Economics of Risky Behaviors, 2015 Western Economic Association Annual Meeting, 2015 Society of Labor Economics Meetings, and the 2015 NBER SI Economics of Crime and Labor Studies Groups. We would also like to thank Tena Ward and Ed Norman from the North Carolina Childhood Lead Poisoning Prevention Program; Julia Rush of the Mecklenburg County Sheriff's Department; Andy Baxter from Charlotte-Mecklenburg Schools; David Jacobs of National Center for Healthy Housing and Diana Adams of LeadSafe Charlotte. This research was partially supported by the Australian Research Council Centre of Excellence for Children and Families over the Life Course (project number CE140100027). The views expressed herein are those of the authors and are not necessarily those of the Australian Research Council.

Lead (Pb) pollution is a pervasive threat to childhood health and development since it is associated with substantial cognitive and behavioral impairments. Despite a dramatic decline in the prevalence of lead due to the prohibition of leaded gasoline, lead exposure is still widely recognized as a major public health issue. Jacobs et al. (2002) estimate that one out of every four homes in the United States contains a significant lead paint hazard. In 2000, the World Health Organization estimated that 40 percent of children under five have levels of exposure associated with neurological damage with 97 percent of these children living in developing countries (Prüss-Üstün et al., 2004). As is the case with other environmental hazards, lead is heavily concentrated in disadvantaged communities and therefore contributes to the intergenerational transmission of inequality through its impact on early-life health (Aizer and Currie, 2014).

Given the large body of evidence connecting childhood lead exposure to cognitive and behavioral deficiencies,² the U.S. Center for Disease Control (CDC) recommends blood lead testing for children around one and two years of age and a case management approach for children whose detected blood lead levels (BLLs) exceed an alert threshold. To reduce childhood exposure and mitigate long-term damage, public health officials implement a combination of actions to both remove lead exposure through information and remediation as well as provide additional health and public assistance benefits for lead-poisoned children.

We merge blood lead surveillance data, public school records, and criminal arrest records at the individual level to evaluate the long-term impact of elevated BLL interventions on school performance and adolescent behavior in Charlotte, North Carolina.³ Similar to that of many other state and local health departments, the public health response in North Carolina is based on CDC guidelines. Two consecutive test results over an alert threshold of 10 micrograms of lead per deciliter of blood ($\mu\text{g}/\text{dL}$) triggers an elevated BLL intervention. Individuals exceeding this threshold for only one BLL test result are not eligible for the elevated BLL intervention.

To identify a causal impact of elevated BLL interventions, we compare a range of behavioral and educational outcomes between our intervention-eligible group—those with an initial and confirmatory BLL test over the alert threshold of $10\mu\text{g}/\text{dL}$ —and a control group. We choose a control group with similar initial BLL test results, but whose confirmatory test falls just below the alert threshold (between 5 and $10\mu\text{g}/\text{dL}$). This group is very similar to those

²EPA (2013) provides an extensive review of hundreds of studies investigating the effects of lead from epidemiology, toxicology, public health, neuroscience, and other medical disciplines. Effects are found across different measures of cognition and academic performance such as: IQ tests (Schnaas et al., 2006; Lanphear et al., 2005; Ris et al., 2004; Canfield et al., 2003; Bellinger et al., 1992), primary school assessments (Aizer et al., 2016; Rau et al., 2013; McLaine et al., 2013; Zhang et al., 2013; Reyes, 2011; Chandramouli et al., 2009; Miranda et al., 2009; Nilsson, 2009; Miranda et al., 2007), high school graduation (Nilsson, 2009; Fergusson et al., 1997; Needleman et al., 1990), and even lower adult earnings (Nilsson, 2009). Early-life lead exposure also impacts externalizing behaviors such as attention, impulsivity, and hyperactivity in young children (Froehlich et al., 2009; Chen et al., 2007); increased delinquent and antisocial activity and higher rates of arrest (Aizer and Currie, 2017; Reyes, 2015; Wright et al., 2008; Fergusson et al., 2008; Needleman et al., 2002; Dietrich et al., 2001; Needleman et al., 1996).

³Charlotte contains the eighteenth largest school district and is representative of other large urban areas in the United States.

eligible for intervention across a wide range of observable characteristics. In our setting, a regression discontinuity model is not ideal due to a small number of observations around the threshold and a growing intensity of the intervention as BLL results increase from the threshold.⁴ Therefore, we take advantage of the well-established negative association between lead exposure and childhood development. In this context, an impact of higher levels of exposure among our treatment group will bias our estimates of the elevated BLL intervention towards finding no effect. However, our preferred treatment and control groups are balanced across observable characteristics and those in the treatment group do not appear to live in more risky environments as measured by prior parcel-level BLL test results. These factors alleviate concerns about a large understatement of the intervention benefits in our sample caused by any higher risks of exposure among the treatment-eligible individuals.⁵

All cases with two BLL tests exceeding the alert threshold ($10\mu\text{g}/\text{dL}$) trigger eligibility for an intervention which includes the following actions: education for caregivers (which includes nutritional advice and information about reducing exposure in the home); a voluntary home environment investigation; and a referral to lead remediation services. A more intensive intervention can be triggered by tests over $15\mu\text{g}/\text{dL}$ or $20\mu\text{g}/\text{dL}$. In addition to educating caregivers and providing a referral to remediation services, the intensive intervention typically includes: a mandatory home environment investigation; nutritional assessment; medical evaluation; developmental assessment; and a referral to the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC).

We estimate a substantial decrease in antisocial behavior among individuals whose BLL test results trigger eligibility for an intervention. Relative to our control group, we find a 0.184 standard deviation decrease in antisocial behavior for adolescents using a summary index. We also estimate a marginally significant 0.117 increase in primary and middle school educational performance among children eligible for an intervention that is administered prior to school entry.⁶ These intention-to-treat estimates are large in magnitude.⁷ In fact, our results suggest that the effects of high levels of exposure on antisocial behavior can largely be reversed by the intervention—children who test twice over the alert threshold exhibit similar outcomes as children with lower levels of exposure ($\text{BLL} < 5\mu\text{g}/\text{dL}$).

⁴We present results from several different regression discontinuity designs and provide plots of outcomes by BLL values in the Appendix.

⁵Similarities between our treatment and control groups may not be that surprising given well-known issues with blood lead tests in accurately measuring exposure risk (ATSDR, 2007; Kemper et al., 2005; CDC, 1997). First, blood tests are better suited to detect contemporaneous shocks rather than cumulative exposure because lead has a short half-life (approx. 30 days) in the blood stream. Second, blood is often drawn through a capillary (finger-prick) sample which carries a high risk of contamination.

⁶For educational and behavioral outcomes we pool a large set of primary outcomes into two summary indexes to limit multiple hypothesis testing concerns previously identified among evaluations of early-life interventions (Anderson, 2008).

⁷We estimate the effects of intervention eligibility (intention-to-treat effects) because we do not have information on intervention compliance.

Our study offers two primary contributions. First, we provide novel estimates of the long-term impact of the standard public health response to elevated BLLs among young children in the United States. Since the CDC lowered the alert threshold to 10 μ g/dL and published new recommendations in 1991, millions of children in the United States would have been eligible for the early-life health and environmental treatments following results of elevated blood lead levels in states which follow the CDC recommended response.⁸ Despite this large-scale public health response to lead-poisoned children, no previous studies evaluate whether there are long-term behavioral or educational benefits associated with these environmental and health interventions.

Second, this paper contributes to a growing literature evaluating the causal impact of early-childhood health interventions on long-term cognitive and behavioral outcomes (Cunha and Heckman, 2008; Currie and Almond, 2011). Recent research suggests that early health and education interventions can yield large long-term benefits.⁹ The Carolina Abecedarian Project—which provided a package of treatments focused on social, emotional, and cognitive development to disadvantaged children from birth through age five—has been associated with increases in educational attainment, reductions in criminal activity, and improved adult health (Barnett and Masse, 2007; Anderson, 2008; Campbell et al., 2014). Many other early-life interventions have also proven effective, such as those administering increased medical care at birth (Bharadwaj et al., 2013); nutritional supplementation for pregnant women and young children (Hoynes et al., 2011); nurse home visit programs (Olds et al., 1999, 2007); and high-quality preschool programs such as Perry Preschool and Head Start (Currie and Almond, 2011; Heckman et al., 2013; Bitler et al., 2014; Conti et al., 2015). The elevated BLL intervention is unique to this literature because it has been widely applied as a public health response to an environmental toxin.

The primary goal of intervention following a confirmed elevated blood lead level is to prevent further exposure and to reduce lead levels in affected children. Two primary channels emerge through which intervention affects antisocial behavior and cognitive outcomes. First, intervention may dramatically reduce the amount of continued childhood exposure to the dangerous neurotoxin by directly reducing exposure risks within the home environment.¹⁰ Second, long-term benefits may occur through improvements in early-life health unrelated to any changes

⁸Since the CDC began collecting national statistics on blood lead surveillance in 1997, nearly one million children were confirmed to have elevated BLLs (BLL > 10 μ g/dL) (surveillance statistics obtained from <http://www.cdc.gov/nceh/lead/data/national.htm> [accessed Jan 24 2015]). Projecting these testing rates and results back to 1991 and assuming that states follow the CDC recommended procedures implies millions of confirmed elevated BLL cases that trigger intervention. While millions may have been eligible, we do not have information on compliance rates with interventions in NC or in other states.

⁹See Currie and Almond (2011) for a recent review.

¹⁰Benefits from reductions in environmental lead levels are expected given several recent studies showing evidence of a causal relationship between exposure and long-term outcomes (Aizer and Currie, 2017; Reyes, 2015; Clay et al., 2014; Grønqvist et al., 2014; Rau et al., 2013; Ferrie et al., 2012; Reyes, 2011; Nilsson, 2009; Troesken, 2008; Reyes, 2007).

in lead exposure.¹¹ We cannot separately identify these two mechanisms or estimate the effects of specific elements of these elevated BLL intervention packages.¹² However, we do present evidence suggesting that both mechanisms contribute to long-term benefits. We find that households in our treatment group that are more likely to have reduced exposure, such as those with children who experience an immediate and sharp decline in post-intervention BLL test results, experience larger benefits. On the other hand, we estimate large effects for individuals eligible for treatments not directly addressing exposure risk, suggesting that long-term benefits should be at least partially attributed to general improvements in early-childhood health.

While further research is needed to investigate the mechanisms by which individuals benefit from elevated BLL interventions, cognitive and behavioral effects associated with the standard intervention package are still relevant in evaluating current public health policy. Public health organizations now state that no BLL should be considered “safe” and have recommended lowering the threshold to identify additional children at risk for health and developmental problems caused by exposure to lead (Budtz-Jørgensen et al., 2013; CDC, 2012).¹³ Applying similar interventions at lower BLL thresholds may yield a large return on investment considering the magnitude of our estimates and the large returns previously associated with other early childhood interventions.¹⁴

The remainder of the paper is structured as follows: Section 2 describes the early-life interventions triggered by elevated BLLs in Charlotte, NC. Section 3 describes our data and characterizes our intervention and control groups with summary statistics. Section 4 outlines our empirical strategy to identify causal effects of intervention. Section 5 presents and discusses estimated effects on a variety of educational and behavioral outcomes, and Section 6 investigates the mechanisms driving our main results. Finally, Section 7 provides a simple cost-benefit analysis and Section 8 provides some concluding remarks. An Appendix contains detailed descriptions of data sources and includes additional analysis.

¹¹The elevated BLL intervention package includes treatments previously demonstrated to impact later-life outcomes such as: visits from health workers; increased medical care; nutritional assessments and dietary modifications; and referral to the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC). Prior research documents long-term benefits from programs similar to each of these elements: increased medical care at birth (such as those triggered by Very Low Birth Weight evaluated by Bharadwaj et al. (2013)); increased access to medical professionals (e.g. the Nurse-Family Partnership evaluated by Olds et al. (2007)); improved early-life nutrition and increased access to public assistance programs (Hoynes et al., 2011, 2012); high-quality early childcare and preschool programs which focus on these social and cognitive developmental processes (e.g. Abecedarian, Perry Preschool, and Head Start).

¹²The majority of evaluations of other early-life interventions also estimate effects for an intervention package containing several components. For example, the original Abecedarian intervention combined early education with a nutritional and health component (Campbell et al., 2014); Bharadwaj et al. (2013) find long-term effects from a “bundle of medical interventions” triggered by a very low birth weight threshold.

¹³The NC Childhood Lead Poisoning Prevention Program of the Children’s Environmental Health Branch currently provides more information about nutrition and key sources of exposure for children testing over 5µg/dL.

¹⁴Cost benefit analyses of early-life intervention programs find a 4 to 1 return for Abecedarian (Masse and Barnett, 2002) and a 7 to 1 return associated with Perry Preschool (Karoly et al., 1998).

2 Description of Public Health Interventions Triggered by Elevated Blood Lead Levels

The U.S. Center for Disease Control and Prevention (CDC) currently funds the development of state and local childhood lead poisoning prevention programs and surveillance activities with the following objectives: to screen infants and children for elevated blood lead levels; to refer lead-poisoned infants and children to medical and environmental interventions; to educate healthcare providers about childhood lead poisoning; and to implement preventative measures to reduce childhood exposure (Meyer et al., 2003). In 1991, the CDC defined a blood lead level of $10\mu\text{g}/\text{dL}$ as the “level of concern” and recommended the provision of specific medical and environmental services from public health agencies following blood lead tests exceeding this threshold (CDC, 1991).¹⁵

The NC Childhood Lead Poisoning Prevention Program of the Children’s Environmental Health Branch bases intervention policies and procedures on CDC recommendations.¹⁶ The standard experience of a child that may be at risk for lead exposure is through an initial BLL test as part of a regular scheduled doctor’s visit between the ages of one and two. In our data, we see a large number of visits clustered around 12, 18 and 24 months of age consistent with this experience. The timing of confirmatory testing is recommended to occur within one month for BLL values $> 20\mu\text{g}/\text{dL}$ and a typical patient usually returns to the same health provider as the initial test and gets re-tested about 4 months later.

If an initial test indicates a blood lead level greater than $10\mu\text{g}/\text{dL}$, a confirmation test is required within six months.¹⁷ If a second consecutive test indicates a blood lead level greater than $10\mu\text{g}/\text{dL}$, a set of interventions is implemented based on the level of lead detected.¹⁸ Figure 1 documents CDC recommendations as of 2002. Based on conversations with health workers in Mecklenburg County, NC, these CDC recommendations constituted public health policy in

¹⁵The intervention level was $25\mu\text{g}/\text{dL}$ between 1985 and 1991; $30\mu\text{g}/\text{dL}$ between 1975 and 1985; and $40\mu\text{g}/\text{dL}$ between 1970 and 1975 (CDC, 1991).

¹⁶The state of North Carolina recommends blood lead tests for all children at age 12 months and again at age 24 months. In practice, the children screened for lead is limited to those individuals who live in neighborhoods with older homes (pre 1978) and when a child’s parents answer “yes” or “don’t know” to any questions on the CDC lead risk exposure questionnaire. The state of NC also requires lead testing for individuals participating in the Medicaid or WIC programs.

¹⁷Confirmatory tests after 6 months are not valid and thus considered a second initial test by the state.

¹⁸The initial test is usually based on capillary specimens typically obtained by the a finger prick where the recommended procedure for a follow-up test is through venous blood draw, which is less likely to be contaminated. Surprisingly, the blood Lead Surveillance data indicate that approximately one-third of follow-up tests are venous during our sample period. The lack of compliance with this aspect of the CDC recommendations is potentially due to local health workers preferring the less-invasive capillary specimen method. We find no systematic differences across the treatment and control in the type of the confirmatory test and find that the initial lead value is not predictive of the second test type. These results indicate that the variation in confirmatory test type is likely due to resources available at the testing clinics and local health worker preferences.

Charlotte back to 1991.¹⁹

The set of interventions for our entire sample of children with two consecutive tests over 10µg/dL include the following: provision of nutritional and environmental information; a referral to WIC for families not already participating; an environmental history interview to identify sources of lead; and a referral to remediation programs for cases identified as high lead risk in the home. Tests over 15µg/dL or 20µg/dL can initiate a more intensive intervention in which children also receive the following treatments: a mandatory home environmental investigation; a medical evaluation; and a detailed nutritional assessment. We test for heterogeneous intervention effects for children with BLLs over these thresholds. According to conversations with individuals from the North Carolina Childhood Lead Poisoning Prevention Program, interventions are only substantially different at the 20µg/dL threshold in practice. This increase in intensity of intervention at the 20µg/dL threshold is evident in Figure 1 which emphasizes more direct medical and remediation action and is also supported by our estimates.

The formal protocol for the standard intervention includes first taking a medical history regarding any symptoms or developmental problems along with previous blood lead measurements and family history of lead poisoning. The healthcare provider then performs an environmental history interview during which family members are asked about the age, condition, and ongoing remodeling or repainting of a child's primary residence as well as other places where the child spends time (including secondary homes and childcare centers). The healthcare provider then determines whether a child is being exposed to lead-based paint hazards at any or all of these places. The environmental history also includes an inquiry about other sources of potential lead exposure.²⁰

Based on the environmental history interview or a confirmatory test over 20µg/dL, a professional lead remediation team conducts a lead inspection at the child's home. This inspection leads to a determination of the home being lead-safe or in need of lead remediation. The provision of lead remediation services involves the removal of lead contaminants, which usually requires the replacement of windows and doors and the repainting of interior/exterior walls. During our sample time period, lead remediation was primarily funded through local government agencies, HUD based lead remediation grants, nonprofits and privately. The cost for lead remediation is not trivial with the average price of these repairs totaling \$7,291.²¹

¹⁹We have found no evidence of any changes in policy preceding 2002 when the CDC recommendations were published in the NC Childhood Lead Poisoning Prevention Program lead testing manual. Since the mid 2000s, procedures have changed slightly to include the provision of nutritional and environmental information for individuals testing over 5µg/dL. However, during the time period when our sample was tested for lead (1990-2000), the 5µg/dL threshold did not trigger any policy interventions.

²⁰Some additional sources of lead include Vinyl miniblinds manufactured prior to 1996, soil and dust which is primarily contaminated by previous existence of lead paint, leaded gasoline or pipes, as well as toys and pottery from overseas.

²¹This estimated cost is based on cost data from LeadSafe Charlotte, which began operations in 1998 and was funded by HUD to remediate lead from homes in Charlotte.

Since lead levels in the body are the result of a combination of lead exposure and the body's absorption of lead into the brain, nutrition can mitigate the effects of lead exposure. While the effectiveness of nutritional interventions is not established, research suggests that deficiencies in iron, calcium, protein, and zinc are related to BLLs and potentially increase vulnerability to negative effects of lead (CDC, 1991). A nutritional assessment includes taking a diet history with a focus on the intake of iron-, vitamin C-, calcium-, and zinc-rich foods. The nutritional information is also used to assess the ingestion of non-food items as well as water sources that contain lead for the family. The healthcare provider inquires into participation in WIC or the Supplemental Nutrition Assistance Program (SNAP or "food stamp") and refers the family to these programs if they are not currently participating. For children with a confirmatory test over 20µg/dL, a medical examination is conducted with particular attention to a child's psychosocial and language development. In cases of developmental delays, a standardized developmental screening test is recommended, which offers referrals to an appropriate agency for further assessment.

3 Data

We merge blood lead surveillance data, public school records, and criminal arrest records at the individual level for children born between 1990 and 1997 in Charlotte-Mecklenberg County, NC.²² Blood lead surveillance data are maintained by the NC Childhood Lead Poisoning Prevention Program of the Children's Environmental Health Branch.²³ This dataset includes BLL test results, which allow us to determine which children were eligible for various lead policy interventions due to two tests with BLLs of 10µg/dL or above.²⁴

We match individual children who receive blood lead tests to two additional databases in order to examine the impact of elevated BLL interventions on educational and behavioral outcomes. First we match BLL test results to administrative records from Charlotte-Mecklenburg Schools (CMS) that span kindergarten through 12th grade and the school years 1998-1999 through 2010-2011.²⁵ Specifically, we incorporate student demographics on race and home address,

²²We restrict our sample to individuals born in 1997 or earlier to allow all individuals to reach age 16 by 2013.

²³North Carolina requires all children participating in Medicaid or the Special Nutrition Program for Women, Infants, and Children (WIC) to be screened for lead at one or two years of age. Other children are screened if a parent responds "yes" or "don't know" to any of the questions on a CDC Lead Risk Assessment Questionnaire. The North Carolina Blood Lead Surveillance Group estimates that it screened between 21.9 and 30.4% of children one and two years of age from 1995 through 1998 and we expect screening rates were similar during our analysis period Miranda et al. (2007).

²⁴These data also include a child's name, gender, birth date, test date, BLL, and home address.

²⁵We are able to match 74 percent of individuals with two tests and one test >10µg/dL in the blood lead surveillance data to a student record in CMS. We do not find any statistically significant differences in the ability to match blood lead test data to CMS school records (which is required for an individual to be included in our estimation sample) and report a match rate comparison in the Appendix (Table A5).

yearly end-of-grade (EOG) test scores for grades 3 through 8 in math and reading,²⁶ number of days absent, days suspended from school, and the number of incidents of school crime.²⁷

To examine adult criminal outcomes, we match our lead database to a registry of all-adult (defined in North Carolina as age 16 and above) arrests in Mecklenburg County from 2006 to 2013.²⁸ The arrest data include information on the number and nature of charges as well as the date of arrest. These data allow us to observe adult criminality regardless of whether a child later transferred or dropped out of school, the main limitation is that it only includes crimes committed within Mecklenburg County. While we cannot observe whether individuals are more or less likely to leave Mecklenburg County following school, we do not find any statistically significant difference in the probability of attending public high school in the county which alleviates any concerns about differential attrition/mobility between our treatment and control groups.²⁹

We draw on two additional databases to control for parental and housing factors, which may influence outcomes. The first data are the population of birth certificate records from the state of North Carolina from 1990-1997 from which we obtain birth weight and years of parental education.³⁰ The second database is county assessor's data for all parcels. Property data can be matched to lead test results based on home address. We augment this parcel data with building permits for all home renovations between 1995 and 2012. This database allows us to incorporate information on housing stock and neighborhoods, directly accounting for some degree of home maintenance that may be correlated with lead exposure. This database on parcels allows us to generate variables for prior home renovations, age, and type of housing structure.³¹ We match our sample to these two data sets but do not require a match for a observation to be included in our estimation sample. Instead, we create dichotomous variables indicating a non-match across the birth record and parcel databases and assigned missing variables a value of zero.³²

Tables 1 and 2 display summary statistics for our intervention group and control group (defined

²⁶Test scores are standardized at the state level by grade and year.

²⁷According to NC State Statute 115C – 288(g), any incident at school involving any violent or threats of violent behavior, property damage, theft or drug possession must officially be reported to the NC school crimes division. This statute ensures that this measure of school crime is consistently reported across schools and cannot be treated differently based on school administrators.

²⁸We use first name, last name and date of birth to link individuals across the two data sources. Details are provided in the Appendix.

²⁹These results are presented in the Appendix (Table A5).

³⁰We are able to match approximately 54 percent of birth records to our lead database. We do not limit our estimation sample by a match to this database; we create indicator variables for any individuals we are unable to match to the birth record database. Even though this match rate is somewhat lower than our other databases, the variables from this database are simply used as control variables and we later show that this match rate is unrelated to our lead policy intervention group.

³¹The lead database is matched to parcel records 86 percent of the time with differences primarily a result of incomplete home address information.

³²We also do not find significant differences in whether we are able to match individuals in our estimation sample to the birth record or parcel information databases and report this comparison in the Appendix (Table A5).

in Section 4) after merging all data and limiting our analysis to individuals born prior to 1998.³³ Individual attributes are similar between the two groups in our estimation sample (Table 1), yet the intervention group has substantially better education and behavioral outcomes (Table 2). We further explore these differences through a regression analysis discussed in the following section.

4 Empirical Framework

In order to assess the impact of the early-life interventions triggered by elevated BLLs, we estimate the following model for all individuals who have an initial BLL test above 10 μ g/dL, return for a second test, and can be matched to CMS public school records:

$$Y_i = \alpha \text{Intervention}_i + \mathbf{X}_i \beta + \epsilon_i \quad (1)$$

where Y_i is an outcome for individual i and \mathbf{X}_i includes a wide range of controls.³⁴ Each outcome is regressed on an indicator, Intervention_i , for whether child i received two consecutive tests over the intervention threshold of 10 μ g/dL. Since the presence of lead paint is heavily concentrated in older residential neighborhoods, standard errors are clustered at the Census Block Group (CBG) level.³⁵

Our primary results focus on intervention effects for two summary index outcomes: educational performance and adolescent antisocial behavior. We follow the methodology for creating a summary index as outlined in Anderson (2008) in a re-evaluation of several early childhood intervention programs.³⁶ Besides dealing with concerns about multiple hypothesis testing, a summary index can be potentially more powerful than individual-level tests due to random error in outcome measures. The antisocial behavior index includes measures of absences and

³³We provide a table describing all variables used and their sources in the Appendix (Table A1). We also present summary statistics for the entire population after merging all data (Tables A2 and A3). In general, we observe lower educational and behavioral outcomes for children who receive a blood lead test compared to untested children and worse outcomes for those with high detected BLLs relative to those with minimal BLLs. Lead tests and higher test results are more likely among children living in older homes, lower income neighborhoods, and with less parental education.

³⁴We include indicators for gender, race/ethnicity, birth year, single family home and home built pre 1978 as well as continuous variables for age at blood test, birth weight, parental education level, the average previous lead test results associated with the residential address listed, as well as an index measure of disadvantage derived from census block group variables (based on address at first lead test). We calculate this summary neighborhood measure from an unweighted z-score sum of the percent of households without a high school graduate, the CBG poverty rate, the fraction of single female headed households, and the CBG population density.

³⁵There are 151 CBGs in our primary analysis. Given the downward bias detected when the number of observations differs across groups or for other forms of cluster heterogeneity in Carter et al. (2017), we also calculate the *effective number of clusters* around 30 for our regressions. This level is not associated with substantial bias in Carter et al. (2017).

³⁶The steps to calculate the summary index are outlined in detail in Anderson (2008). We also provide a description of the steps in the Appendix.

number of days suspended (6th through 10th grade); school reported crimes, and adolescent criminal arrests from the age of 16 through 18.³⁷ The educational performance index includes 3rd through 8th grade math and reading test score results as well as grade retention between 1st and 9th grade.³⁸ We also estimate and present results separately for individual outcomes used in the summary indexes.

Throughout the empirical analysis, we estimate Eq. (1) restricting our sample to individuals with an initial BLL test of 10 μ g/dL or greater. Our primary control group includes individuals who have one test over the alert threshold of 10 μ g/dL and the confirmatory test within six months between 5 and 9 μ g/dL. Figure 2 illustrates the various combinations of BLL values among all individuals with at least two tests with the larger font numbers highlighting the individuals that populate our treatment and control groups.³⁹

Our identification strategy relies on the fact that our treatment and control groups both receive a BLL test with an elevated initial value and follow-up public health recommendations for re-testing BLL. This criteria should address most issues of selection on testing as well as parental and environmental differences that may impact the ability or desire to re-test. Since the test results are, on average, higher for those eligible for the intervention package, this group may experience a more dangerous level of underlying lead exposure which, based on previous literature, is associated with larger education and behavioral deficits. Figures 3 and 4 clearly depict more negative outcomes for reading/math test scores, repeat grades in school, suspensions, absences and crime associated with higher BLL values.⁴⁰

To assess whether intervention is unrelated to unobserved determinants of cognitive and behavioral outcomes, we compare observable characteristics (including measures of parental quality, health at birth, housing quality, and neighborhood quality) across the intervention and control groups. Despite large and statistically significant differences between mean outcomes in Table 2, we find no significant differences among observable characteristics between our intervention and control groups in Table 1. The small differences in individual attributes between the intervention and control group is formally investigated in a balance test presented on the bottom of Table 1—we cannot reject that all variables are jointly equal to zero.⁴¹ Notably, we do not find a statistically significant difference in prior BLLs tied to the parcel addresses of our treatment and control groups, suggesting that exposure risks are similar across the two

³⁷We treat the absences as coming from truancy, a behavioral outcome, but note that absences could also be due to health problems.

³⁸We limit our analysis to school outcomes through 10th grade because our public school records are available only through the 2010-2011 school year and we have very few cohorts in 11th or 12th grade by 2010. Criminal arrest data is available for an additional 2.5 years (through 2013) allowing us to measure arrests between 16 and 18 years of age for many of the children receiving lead tests since 1992.

³⁹ We plot the distribution of BLL values for all first and second test values in the Appendix (Figure A8).

⁴⁰In the Appendix, Table A2 presents a similar pattern for the full population of CMS students with BLL tests born between 1990 and 1997.

⁴¹Results from the balance test specification are presented in the Appendix (Table A4).

groups.⁴² This result points towards measurement error as an important source of variation in treatment eligibility within our estimation sample.⁴³

Throughout our analysis we refer to our estimates as intervention effects, but our estimated effects represent a combination of several responses to intervention. First, since we do not directly observe participation in any intervention programs, our estimated effects are intention-to-treat (or “ITT”) treatment effects which represent a combination of the direct impact of intervention on outcomes and the probability of compliance with the intervention.⁴⁴ Second, the estimated impact includes the role of parental or other inputs that react to a confirmed elevated BLL. For example, intervention could directly impact child nutrition and the level of lead in the home environment but also impact the amount of care and attention provided by a parent. While decomposing the various components of this total effect would be extremely useful in designing early childhood intervention programs, our estimated intervention effect is more relevant for the evaluation of the CDC-recommended public health response to elevated BLLs. The effect of the policy will always include direct benefits of intervention, potential non-compliance, and any indirect benefits from family or community responses to intervention.

5 Results

After a second test confirms an elevated BLL, the NC Department of Health requires the implementation of the interventions recommended by the CDC (as listed in Figure 1). The CDC recommends testing until an individual with elevated levels tests below the alert threshold of $10\mu\text{g}/\text{dL}$. To assess whether individuals comply with intervention after an elevated BLL is confirmed, we estimate the effect of intervention on several measures of continued testing. Columns (1) through (3) of Table 3 demonstrate that compared with the control groups, those with confirmed elevated BLLs are 44 percentage points more likely to have a third test within 100 days of the confirmatory BLL test result, have twice as many overall tests, and respond quickly following a second elevated test by obtaining a third test within approximately three months. Overall, 79 percent of individuals in our intervention group continue testing until their $\text{BLL} \leq 10\mu\text{g}/\text{dL}$ (as depicted in Figure 5). While these results provide some confidence that, on average, interventions are administered to children who are supposed to receive them according to local health department policy, all of our estimates remain intention-to-treat estimates since

⁴²We also compare outcomes across those living in the same parcels *prior* to the children in our estimation sample using the same empirical specification (Eq. 1) and report these results in Table 6. Despite our treatment group having higher detected BLLs, we do not find evidence from these specifications that these higher BLLs are driven by differences in the home environment.

⁴³Blood testing is a noisy measure of exposure for two reasons: 1) a short half-life of lead in blood (30 days) and 2) a high risk of contamination during testing procedures that utilize capillary sampling (ATSDR, 2007; Kemper et al., 2005; CDC, 1997).

⁴⁴It is possible that some families refuse any intervention after two consecutive tests over the alert threshold. These families would be “treated” in our framework since we do not observe implementation.

we do not have data indicating participation in the components of the intervention package. The first panel of Table 4 estimates Eq. (1) for our education summary indexes and for individual outcomes grouped by different grade levels. Combining math and reading test scores between the 3rd and 8th grade as well as grade retention outcomes between the 1st and 9th grade into a summary index, we estimate a marginally significant 0.117 standard deviation increase in educational performance associated with the elevated BLL intervention. While the majority of our test score estimates are imprecise, they are at least consistent with benefits from intervention in direction and magnitude.

The second panel of Table 4 reports a large and significant decline in antisocial behavior associated with elevated BLL intervention. Relative to the control group, we estimate a 0.184 standard deviation decrease in our antisocial behavior summary index associated with intervention. This represents a very large drop from the average index value of 0.10 for the control group. The pattern of estimates across individual outcomes of suspensions, absences, school crimes, and criminal arrests reported in Table 4 consistently demonstrates improvements associated with intervention.

Overall, the pattern of our estimated effects are consistent with recent work suggesting that effects from early-childhood interventions which boost non-cognitive skills do not fade out over time (Heckman et al., 2013). The magnitude of the difference between control and treatment group outcomes grow with later outcomes—we find larger effects for the later test score results compared to primary school test scores and the largest impact on secondary school behavior outcomes.

In Figure 6, we estimate treatment effects relative to six different control groups defined by a range of the average detected BLLs. Note that the control groups in these specifications are no longer restricted only to individuals with an initial test above $10\mu\text{g/dL}$. This plot accomplishes two objectives: First, the plot demonstrates that our estimated treatment effects are consistent with the idea of worse outcomes for individuals with higher BLL values since we only find sizeable and significant effects when we compare our treatment group to control groups with higher BLL values (5 and above). Second, the plot provides an interesting framing of our estimated treatment effects in that we do not find statistically significant differences between our intervention eligible group and children detected to have BLL levels below $5\mu\text{g/dL}$. We find that the elevated BLL intervention can largely mitigate the education and behavioral deficiencies associated with higher levels of exposure.

In our context, higher levels of underlying exposure among our treatment group would lead to a downward-biased estimate of the benefits from the elevated BLL policies. We provide a number of results that show limited evidence of such a downward bias. First, we do not find strong evidence of a large difference in exposure risk when comparing average BLL test results within treatment and control parcels in years prior to the testing of our treatment

and control individuals (Table 2). We also do not find evidence that observable characteristics correlated with higher levels of disadvantage (e.g. low birth weight, low maternal education, high neighborhood poverty) are consistently worse among the intervention group relative to the control group.⁴⁵

The lack of any statistical difference in demographic and parcel characteristics correlated with higher lead exposure risks suggests that, in this setting, a large part of the variation in treatment eligibility could be explained by idiosyncratic variation in measuring exposure through blood lead testing. Specifically, several characteristics of blood lead testing support measurement error as an important source of variation in test results. First, BLL levels are influenced by the relationship between date of exposure (which is usually unknown to the family) and the date of testing with only a month of passed time generating over a 50 percent decrease in the BLL due to a short half-life of lead in blood (30 days). Second, capillary sampling (a “finger-prick” method) is the most common type of test for both initial and confirmatory tests in Charlotte during our time period of analysis and is known to have a high contamination risk relative to alternative testing procedures.⁴⁶

We present additional results in an Appendix to assess whether our estimates are robust to alternative specifications. These exercises include: the influence on various sets of controls on estimated effects (Table A6); specifications which include other control variables than our baseline set described in Section 4 (Table A7); and estimates using regression discontinuity designs (Table A8). Given the size of our estimation sample, estimates from the regression discontinuity specifications or those including a more extensive set of controls are usually similar in direction and magnitude but are less precise.⁴⁷

We also evaluate whether elevated BLL intervention impacts other children in the household by matching intervention and control individuals to their siblings. Among a small number of sibling observations, we find patterns consistent with there being an effect of intervention for the household with the benefits concentrated among younger siblings (Table A9). To the extent interventions reduce levels of dangerous lead exposure, we expect larger effects for younger siblings since older siblings would already be damaged from exposure. We interpret these results cautiously since they are based on few observations and are associated with large

⁴⁵We also do not find that estimated effects increase when including controls for observable characteristics (see Table A6 in the Appendix).

⁴⁶Other non-blood testing procedures, such as measuring lead in children’s teeth, are more accurate but also more expensive and therefore less prevalent. Tooth lead testing is a more accurate measure of cumulative exposure since there is little risk for contamination and due to the fact that the elimination half-life for inorganic lead in bone is approximately 27 years (ATSDR, 2007).

⁴⁷We calculated heterogeneous effects across different demographic groups but these estimates are noisy due to a small number of individuals eligible for treatment in each subsample. For example, the number in the intervention group whom have birth records indicating a parent without a high school degree is 25. Overall, these estimates suggest slightly larger benefits for female children and those with parents who did not graduate from high school. Larger treatment effects for females are also found across evaluations of other early childhood interventions (Anderson, 2008).

standard errors.

6 Mechanisms and Intensity of Treatment

The substantial improvements associated with the elevated BLL interventions likely represent a combination of direct and indirect effects from both the local health department's response and the parental response to lead exposure. Two primary channels emerge through which intervention affects antisocial behavior and cognitive outcomes. First, intervention may dramatically reduce the amount of continued exposure to the dangerous neurotoxin by directly reducing exposure risks within the home environment. Second, long-term benefits may occur through improvements in early-life health unrelated to any changes in lead exposure.

As previously discussed in Section 2 and evident in Figure 1, higher intensity interventions are recommended following confirmatory tests over $15\mu\text{g}/\text{dL}$ and $20\mu\text{g}/\text{dL}$. We explore whether these higher-intensity interventions are associated with larger benefits in the first panel of Table 5. We find substantial benefits among those with a confirmatory test over $20\mu\text{g}/\text{dL}$ and do not detect any additional benefits for those with a confirmatory test between 15 and $20\mu\text{g}/\text{dL}$. Additional effects at the $15\mu\text{g}/\text{dL}$ threshold are not expected in this setting since, according to individuals at the North Carolina Childhood Lead Poisoning Prevention Program, interventions are only substantially different at the $20\mu\text{g}/\text{dL}$ threshold in practice. These results suggest larger benefits from more intensive interventions but are based on a small number of individuals. The larger effects also do not help distinguish between mechanisms since the higher-intensity intervention is associated with more targeted efforts to reduce exposure through mandatory home investigations but is also associated with an increase in medical attention, developmental surveillance, and access to public assistance programs. These results do suggest that the intensity of the local health department's response is potentially an important determinant of long-term benefits and are consistent with prior evaluations of early-life programs. Our point estimates of the lower intensity intervention suggest some benefit, but are not statistically significant which is not surprising given the general lack of power to detect statistically significant effects on subsamples of our treatment group.⁴⁸

Following a second elevated BLL test result, nearly 80 percent of individuals continue to get tested until their BLLs drop below the alert threshold of $10\mu\text{g}/\text{dL}$. While some individuals may test below the threshold due to mean reversion of inaccuracy in testing, many likely have lower BLLs due to some effort to reduce the risk of exposure in the residential environment.

⁴⁸While an important aspect of the lower intensity interventions is parental education about ways to control household exposure, they also provide nutritional information and a referral to remediation services. Thus, these estimated (imprecise) benefits are not inconsistent with previous randomized control trials that do not find large or significant BLL reductions when evaluating parental education and "household dust control" interventions (Campbell et al., 2011; Yeoh et al., 2009; Brown et al., 2006; Jordan et al., 2003; Lanphear et al., 1999).

Reduction in exposure could be due to a parental response to information provided through discussions with health workers following a confirmatory elevated BLL test result or through instructions provided following a home-environment inspection. Reduction could also be due to the provision of remediation services following a home investigation or a referral to available remediation programs.

The most immediate (and expensive) way to reduce environmental exposure within residences identified to contain a lead hazard is through a remediation service. Prior evaluations of household lead remediation programs through randomized controlled trials document significant decreases in levels of household dust (Sandel et al., 2010) and the number of elevated BLL cases (Jones, 2012). If an inquiry or home investigation identifies a potential residence-based hazard for children exceeding the alert threshold, families are typically referred to lead-based paint removal programs. Since 1998, LeadSafe Charlotte, a HUD-funded organization, has provided remediation services to eligible families. While we obtained application and remediation data from this program and are able to match to Charlotte properties, our estimation sample spans birth cohorts between 1990 and 1997, so we cannot match most individuals to remediation services closely following elevated test results. However, we do find a positive association between intervention and whether the parcel was eventually remediated through the LeadSafe Charlotte program in column 4 of Table 3. The magnitude of this coefficient indicates that intervention households were more than three times as likely to have lead remediation as our control group.

To further investigate whether benefits may be due to reductions in levels of exposure, Table 5 compares estimated intervention benefits across individuals in the intervention group who are more likely to have directly addressed lead exposure problems. First, we find larger effects for individuals experiencing a significant drop (more than $5\mu\text{g}/\text{dL}$) between the second and third BLL test. Individuals who experience a sharp drop in BLLs after two consecutive tests over the alert threshold are more likely to have benefited from a reduction in exposure.⁴⁹ We also estimate separate intervention effects for individuals who respond quickly by re-testing within one month following a second test over the alert threshold. The direction of both of these estimates suggests benefits from directly addressing exposure risk.

We also compare outcomes across those living in a “treatment” or control parcel *after* the child in our estimation sample. Table 6 presents results from a specification where individuals living in an intervention parcel after the time of intervention are generally better off along education and behavioral outcomes compared to those living in control households. Also, as discussed earlier, we did not detect any difference in outcomes for individuals matched to the

⁴⁹One may be concerned that with measurement error in BLL tests, an initially high BLL test would be followed by mean reversion for a second test. Since we focus on drops between the 2nd and 3rd tests, the presence of 2 high BLL tests is more indicative of high lead exposure rather than just inaccuracies in testing. Additionally, variation in testing results and subsequent drops in BLL values would only serve to bias the coefficients for Table 5 towards zero.

intervention and control parcels *prior* to BLL testing of our estimation sample. Again, these results mildly suggest that parcels containing a child in the intervention group experience long-term lead exposure reductions.

7 Benefit-Cost Discussion

An important question from a policy perspective remains as to whether the benefits from the elevated BLL intervention outweigh the typical costs of the intervention. While a comprehensive benefit-cost analysis is not feasible in our setting since we cannot yet estimate effects on key outcomes such as employment and earnings, we provide a rough comparison of the typical intervention benefits and costs drawing from previous evaluations of early childhood interventions and estimates of typical costs from administrators of the relevant social service programs in Charlotte.

We estimate the average cost of an elevated BLL intervention in Charlotte at \$5,288. This estimate includes the following components: a doctor's appointment including nutritional assessment/counselling, 3 follow up BLL tests, a home environmental inspection, remediation of lead-based paint hazards by *LeadSafe Charlotte*, and the costs of case management through the Child Development Services Agency. We obtain the information on the costs for each of these components as well as an estimate of the probability the cost is incurred by an individual with two BLL tests over the 10 μ g/dL threshold. This information and the sources for each element are detailed in Table 7.

For an estimate of the intervention benefits, we rely on prior work by Aos et al. (2004) who provide a detailed meta-analysis of the costs and benefits associated with prevention and early intervention programs for youth. Among more than 30 programs reviewed under the category *Early Childhood Education for Low Income 3- and 4-Year Olds*, Aos et al. (2004) calculate an adjusted effect size of 0.08 SD increase in test scores and a 0.162 SD decrease in crime for this category of programs.⁵⁰ These estimated program effects are very similar to our primary estimated effects of 0.117 SD increase in test scores and a 0.184 SD decrease in antisocial behavior (including crime). Given the close proximity of these effect sizes, we draw directly from the estimated benefit calculation from the test score increase and crime decrease in Aos et al. (2004). Aos et al. (2004) estimates the change in expected lifetime earnings associated with a test score improvement of 0.08 SD of \$4,917, and the total social cost savings associated with a 0.162 SD decrease in crime is estimated to be \$4,749.⁵¹ Applying these estimated benefits to our setting implies at least \$9,666 in benefits due to test score improvements and crime reduction

⁵⁰See Table C.1a (pg 16) of Aos et al. (2004).

⁵¹See the *Early Childhood Education for Low Income 3- and 4-Year Olds: Summary of Estimated Benefits and Costs* table on page 94 of Aos et al. (2004) and Appendix D starting on page 33 for a detailed explanation of the benefit calculations for each category.

associated with the elevated BLL intervention. Comparing this to the typical program costs in Table 7 and scaling up the cost estimate by 1.3 to account for the marginal deadweight loss of raising public funds to pay for the intervention (Poterba, 1996), the benefit-cost ratio is 1.4:1. In other words, for each \$1 invested in children with confirmed elevated blood lead levels yields a return of nearly \$1.40 based on our rough (and largely conservative) estimates drawing from prior evaluations interventions for children of similar age and socioeconomic background as those in our estimation sample.⁵²

8 Conclusion

In this first evaluation of the standard public health response to high levels of exposure to environmental lead, we find evidence that interventions can affect long-term educational and behavioral outcomes. We estimate far-reaching decreases in antisocial behaviors (suspensions, school crimes, unexcused absences, and criminal activity) and, to a lesser extent, increases in educational performance. These results support recent evidence that early-life interventions can mitigate and compensate for the deleterious effects of lead.

A massive amount of evidence across multiple disciplines consistently points to a lasting negative impact of lead exposure. In fact, recent studies and media reports suggest that reductions in lead exposure through the prohibition of leaded gasoline may be one of the most important determinants of the decline in crime rates over the past two decades in the United States and other developed nations.⁵³ However, not much is known to what types of programs and policies are effective in addressing these effects. While randomized controlled trials have been used to evaluate other large-scale early childhood interventions (e.g. Head Start), this paper demonstrates that evaluations of interventions related to lead exposure can be conducted using administrative data and by exploiting institutional features (such as testing procedures) to construct a valid counter-factual or control group to evaluate causal effects of intervention.

Although exposure to lead has been reduced in most countries due to the prohibition of leaded gasoline, lead exposure still represents a major public health issue. In the United States, children have continued to be exposed to lead over the last several decades as a result of deteriorating lead paint and contaminated dust within older housing units (Dixon et al., 2009; Gaitens et al., 2009; Levin et al., 2008). The National Survey of Lead and Allergens in Housing estimated that 38 million housing units in the United States (40 percent of all housing units) contained lead-based paint, and approximately 24 million had significant lead-based paint hazards (Jacobs

⁵²This is likely a conservative estimate since it ignores any savings associated with improved health and reduced behavioral problems in school as well as benefits from any spillover effects within classrooms and communities. Moreover, we measure intention-to-treat estimates so the treatment effects would further be scaled up by the rate at which the population complies with the recommended interventions.

⁵³Recent media articles Drum (2013) and Monbiot (2013) highlight this connection based on results from papers by Mielke and Zahran (2012); Nevin (2007); Reyes (2007); Nevin (2000).

et al., 2002). Recognizing the current threat to child health and development in California, a Superior Court judge recently ordered three paint companies to contribute \$1.15 billion to fund the inspection, risk assessment, and hazard abatement of older homes in ten California jurisdictions (Kleinberg, 2014).⁵⁴

Lead exposure is a more pressing public health issue in developing countries where lead in petrol, industrial emissions, paints, ceramics, food and drink cans, water pipes, and traditional medicines is more prevalent. In an evaluation for the World Health Organization, Prüss-Üstün et al. (2004) estimates that 120 million people have blood lead concentrations above 10µg/dL, accounting for an estimated 0.9 percent of the global burden of disease. Prüss-Üstün et al. (2004) also estimates that nearly 10 percent of children under five in the world have blood lead levels greater than 20µg/dL with 99 percent of these children living in developing countries. There is a great deal of evidence that these levels of exposure cause drastic cognitive and behavioral impairment and policies to reduce exposure in developing countries should be of first-order importance.

Until countries and communities make long-run investments in reducing environmental exposure, our results suggest that intervening early is critical to limit the damage from exposure. Our research can be used to inform policymakers considering intervention at lower levels of detected exposure. In 2012, the CDC recognized a lack of evidence for any BLL to be considered “safe” and recommended using a lower threshold to identify children at increased risk for health and developmental problems caused by exposure to lead (CDC, 2012).⁵⁵ It is likely that increasing the frequency and intensity of intervention for lead-exposed children will yield a profound return considering the potential long-term effects of lead on health and human capital.

References

- Aizer, Anna and Currie, Janet.** (2014). ‘The intergenerational transmission of inequality: Maternal disadvantage and health at birth’, *Science* 344(6186), 856–861.
- Aizer, Anna and Currie, Janet.** (2017), *Lead and Juvenile Delinquency: New Evidence from Linked Birth, School and Juvenile Detention Records*, Working Paper w23392, National Bureau of Economic Research.

⁵⁴Judgement was issued for the Plaintiff, the People of the State of California, against Defendants ConAgra Grocery Products Company, NL Industries, Inc. and The Sherwin-Williams Company.

⁵⁵The NC Childhood Lead Poisoning Prevention Program of the Children’s Environmental Health Branch currently provides more information about nutrition and key sources of exposure for children testing over 5µg/dL. The European Food Safety Authority and the World Health Organization have also recently concluded that there is no known safe level of exposure Budtz-Jørgensen et al. (2013).

- Aizer, Anna, Currie, Janet, Simon, Peter and Vivier, Patrick.** (2016), Do Low Levels of Blood Lead Reduce Children's Future Test Scores?, Working Paper w22558, National Bureau of Economic Research.
- Anderson, Michael L.** (2008). 'Multiple inference and gender differences in the effects of early intervention: A reevaluation of the Abecedarian, Perry Preschool, and Early Training Projects', *Journal of the American Statistical Association* 103(484).
- Aos, Steven, Lieb, Roxanne, Mayfield, Jim, Miller, Marna and Pennucci, Annie.** (2004), *Benefits and costs of prevention and early intervention programs for youth*, number Document No. 04-07-3901, Washington State Institute for Public Policy.
- ATSDR.** (2007), *Toxicological profile for lead: Atlanta, GA, USA*, Agency for Toxic Substances and Disease Registry: US Department of Health and Human Services, Public Health Service.
- Barnett, W. Steven and Masse, Leonard N.** (2007). 'Comparative cost-benefit analysis of the Abecedarian program and its policy implications', *Economics of Education Review* 26, 113–125.
- Bellinger, David C., Stiles, Karen M. and Needleman, Herbert L.** (1992). 'Low-Level Lead Exposure, Intelligence and Academic Achievement: A Long-term Follow-up Study', *Pediatrics* 90(6), 855–861. PMID: 1437425.
- Bharadwaj, Prashant, Løken, Katrine Velleesen and Neilson, Christopher.** (2013). 'Early life health interventions and academic achievement', *The American Economic Review* 103(5), 1862–1891.
- Bitler, Marianne P, Hoynes, Hilary W and Domina, Thurston.** (2014), Experimental evidence on distributional effects of head start, Working Paper w20434, National Bureau of Economic Research.
- Brown, Mary Jean, McLaine, Pat, Dixon, Sherry and Simon, Peter.** (2006). 'A randomized, community-based trial of home visiting to reduce blood lead levels in children', *Pediatrics* 117(1), 147–153.
- Budtz-Jørgensen, Esben, Bellinger, David, Lanphear, Bruce and Grandjean, Philippe.** (2013). 'An international pooled analysis for obtaining a benchmark dose for environmental lead exposure in children', *Risk Analysis* 33(3), 450–461.
- Campbell, Carla, Tran, Mary, Gracely, Edward, Starkey, Naomi, Kersten, Hans, Palermo, Peter, Rothman, Nancy, Line, Laura and Hansen-Turton, Tine.** (2011). 'Primary prevention of lead exposure: The Philadelphia lead safe homes study', *Public Health Reports* 126(Suppl 1), 76.

- Campbell, Frances, Conti, Gabriella, Heckman, James J, Moon, Seong Hyeok, Pinto, Rodrigo, Pungello, Elizabeth and Pan, Yi.** (2014). ‘Early childhood investments substantially boost adult health’, *Science* 343(6178), 1478–1485.
- Canfield, Richard L., Henderson, Charles R., Cory-Slechta, Deborah A., Cox, Christopher, Jusko, Todd A. and Lanphear, Bruce P.** (2003). ‘Intellectual Impairment in Children with Blood Lead Concentrations below 10 µg per Deciliter’, *New England Journal of Medicine* 348(16), 1517–1526. PMID: 12700371.
- Carter, Andrew V, Schnepel, Kevin T and Steigerwald, Douglas G.** (2017). ‘Asymptotic behavior of at test robust to cluster heterogeneity’, *Review of Economics and Statistics*, (Forthcoming).
- CDC.** (1991), *Preventing lead poisoning in young children: Recommendations from the Advisory Committee on Childhood Lead Poisoning Prevention: Atlanta, GA, USA*, The Centers for Disease Control (US).
- CDC.** (1997), *Screening Young Children for Lead Poisoning: Guidance for State and Local Public Health Officials: Atlanta, GA, USA*, The Centers for Disease Control (US).
- CDC.** (2012), *Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention: Report to the CDCP by the Advisory Committee on Childhood Lead Poisoning Prevention of the U.S. Centers for Disease Control: Atlanta, GA, USA*, The Centers for Disease Control (US).
- Chandramouli, K, Steer, Colin D, Ellis, Matthew and Emond, Alan M.** (2009). ‘Effects of early childhood lead exposure on academic performance and behaviour of school age children’, *Archives of Disease in Childhood* 94(11), 844–848.
- Chen, Aimin, Cai, Bo, Dietrich, Kim N, Radcliffe, Jerilynn and Rogan, Walter J.** (2007). ‘Lead exposure, IQ, and behavior in urban 5-to 7-year-olds: does lead affect behavior only by lowering IQ?’, *Pediatrics* 119(3), e650–e658.
- Clay, Karen, Troesken, Werner and Haines, Michael.** (2014). ‘Lead and Mortality’, *Review of Economics and Statistics* 96(3), 458–470.
- Conti, Gabriella, Heckman, James and Pinto, Rodrigo.** (2015), *The Effects of Two Influential Early Childhood Interventions on Health and Healthy Behaviors*, Working Paper s21454, National Bureau of Economic Research.
- Cunha, Flavio and Heckman, James J.** (2008). ‘Formulating, identifying and estimating the technology of cognitive and noncognitive skill formation’, *Journal of Human Resources* 43(4), 738–782.

- Currie, Janet and Almond, Douglas.** (2011), Chapter 15 - Human capital development before age five, in **David Card and Orley Ashenfelter.**, ed., 'Handbook of Labor Economics', Vol. Volume 4, Part B, Elsevier, pp. 1315–1486.
- Deming, David J.** (2011). 'Better Schools, Less Crime?', *The Quarterly Journal of Economics* 126(4), 2063–2115.
- Dietrich, Kim N., Douglas, Ris M., Succop, Paul A., Berger, Omer G. and Bornschein, Robert L.** (2001). 'Early exposure to lead and juvenile delinquency', *Neurotoxicology and Teratology* 23(6), 511–518.
- Dixon, Sherry L, Gaitens, Joanna M, Jacobs, David E, Strauss, Warren, Nagaraja, Jyothi, Pivetz, Tim, Wilson, Jonathan W and Ashley, Peter J.** (2009). 'Exposure of US children to residential dust lead, 1999-2004: II. The contribution of lead-contaminated dust to children's blood lead levels.', *Environmental Health Perspectives* 117(3), 468–474.
- Drum, Kevin.** (2013). 'America's Real Criminal Element: Lead', *Mother Jones* (January/February 2013).
- EPA, U.S.** (2013), Integrated Science Assessment for Lead (Final Report), Technical report, U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-10/075F.
- Fergusson, David M, Horwood, L John and Lynskey, Michael T.** (1997). 'Early dentine lead levels and educational outcomes at 18 years', *Journal of Child Psychology and Psychiatry* 38(4), 471–478.
- Fergusson, D M, Boden, J M and Horwood, L J.** (2008). 'Dentine lead levels in childhood and criminal behaviour in late adolescence and early adulthood', *Journal of Epidemiology and Community Health* (1979-) 62(12), 1045–1050.
- Ferrie, Joseph P, Rolf, Karen and Troesken, Werner.** (2012). 'Cognitive disparities, lead plumbing, and water chemistry: Prior exposure to water-borne lead and intelligence test scores among World War Two US Army enlistees', *Economics & Human Biology* 10(1), 98–111.
- Froehlich, Tanya E, Lanphear, Bruce P, Auinger, Peggy, Hornung, Richard, Epstein, Jeffery N, Braun, Joe and Kahn, Robert S.** (2009). 'Association of tobacco and lead exposures with attention-deficit/hyperactivity disorder', *Pediatrics* 124(6), e1054–e1063.
- Gaitens, Joanna M, Dixon, Sherry L, Jacobs, David E, Nagaraja, Jyothi, Strauss, Warren, Wilson, Jonathan W, Ashley, Peter J et al.** (2009). 'Exposure of US children to residential dust lead, 1999–2004: Housing and demographic factors', *Environmental Health Perspectives* 117(3), 461–467.

- Grønqvist, Hans, Nilsson, J. Peter and Robling, Per-Olof.** (2014). ‘Childhood lead exposure and criminal behavior: Lessons from the Swedish phase-out of leaded gasoline’, *Swedish Institute for Social Research Working Paper 9/2014*.
- Heckman, James, Pinto, Rodrigo and Savelyev, Peter.** (2013). ‘Understanding the Mechanisms Through Which an Influential Early Childhood Program Boosted Adult Outcomes’, *The American Economic Review* 103(6), 1–35.
- Hoynes, Hilary, Page, Marianne and Stevens, Ann Huff.** (2011). ‘Can targeted transfers improve birth outcomes?: Evidence from the introduction of the WIC program’, *Journal of Public Economics* 95(7), 813–827.
- Hoynes, Hilary W, Schanzenbach, Diane Whitmore and Almond, Douglas.** (2012), Long run impacts of childhood access to the safety net, Working Paper w18535, National Bureau of Economic Research.
- Jacobs, David E, Clickner, Robert P, Zhou, Joey Y, Viet, Susan M, Marker, David A, Rogers, John W, Zeldin, Darryl C, Broene, Pamela and Friedman, Warren.** (2002). ‘The prevalence of lead-based paint hazards in US housing’, *Environmental health perspectives* 110(10), A599.
- Jones, David J.** (2012). ‘Primary prevention and health outcomes: Treatment of residential lead-based paint hazards and the prevalence of childhood lead poisoning’, *Journal of Urban economics* 71(1), 151–164.
- Jordan, Catherine M, Yust, Becky L, Robison, Leslie L, Hannan, Peter and Deinard, Amos S.** (2003). ‘A randomized trial of education to prevent lead burden in children at high risk for lead exposure: efficacy as measured by blood lead monitoring’, *Environmental Health Perspectives* 111(16), 1947.
- Karoly, Lynn A, Greenwood, Peter W, Everingham, Susan S, Houbé, Jill and Kilburn, M Rebecca.** (1998), *Investing in our children: What we know and don’t know about the costs and benefits of early childhood interventions*, Rand Corporation.
- Kemper, Alex R, Cohn, Lisa M, Fant, Kathryn E, Dombkowski, Kevin J and Hudson, Sharon R.** (2005). ‘Follow-up testing among children with elevated screening blood lead levels’, *JAMA* 293(18), 2232–2237.
- Kleinberg, James.** (2014), Judge Kleinberg’s Order, Final Amended Judgement March 26, 2014 Case Number 1-00-CV-788657, Santa Clara County Superior Court’s Electronic Filing System. Retrieved 16 September 2014.
- Lanphear, Bruce P., Hornung, Richard, Khoury, Jane, Yolton, Kimberly, Baghurst, Peter, Bellinger, David C., Canfield, Richard L., Dietrich, Kim N., Bornschein,**

- Robert, Greene, Tom, Rothenberg, Stephen J., Needleman, Herbert L., Schnaas, Lourdes, Wasserman, Gail, Graziano, Joseph and Roberts, Russell.** (2005). 'Low-Level Environmental Lead Exposure and Children's Intellectual Function: An International Pooled Analysis', *Environmental Health Perspectives* 113(7), 894–899.
- Lanphear, Bruce P, Howard, Cynthia, Eberly, Shirley, Auinger, Peggy, Kolassa, John, Weitzman, Michael, Schaffer, Stanley J and Alexander, Keith.** (1999). 'Primary prevention of childhood lead exposure: a randomized trial of dust control', *Pediatrics* 103(4), 772–777.
- Levin, Ronnie, Brown, Mary Jean, Kashtock, Michael E, Jacobs, David E, Whelan, Elizabeth A, Rodman, Joanne, Schock, Michael R, Padilla, Alma and Sinks, Thomas.** (2008). 'Lead exposures in US children, 2008: implications for prevention', *Environ Health Perspect* 116(10), 1285–1293.
- Masse, Leonard N and Barnett, W Steven.** (2002), *A benefit cost analysis of the Abecedarian early childhood intervention*, National Institute for Early Education Research New Brunswick, NJ.
- McLaine, Pat, Navas-Acien, Ana, Lee, Rebecca, Simon, Peter, Diener-West, Marie and Agnew, Jacqueline.** (2013). 'Elevated blood lead levels and reading readiness at the start of kindergarten', *Pediatrics* 131(6), 1081–1089.
- Meyer, Pamela A, Pivetz, Timothy, Dignam, Timothy A, Homa, David M, Schoonover, Jaime, Brody, Debra et al.** (2003). 'Surveillance for elevated blood lead levels among children-United States, 1997-2001', *Morbidity and Mortality Weekly Report CDC Surveillance Summaries* 52(10).
- Mielke, Howard W. and Zahran, Sammy.** (2012). 'The urban rise and fall of air lead (Pb) and the latent surge and retreat of societal violence', *Environment International* 43, 48–55.
- Miranda, Marie Lynn, Kim, Dohyeong, Galeano, M. Alicia Overstreet, Paul, Christopher J., Hull, Andrew P. and Morgan, S. Philip.** (2007). 'The Relationship between Early Childhood Blood Lead Levels and Performance on End-of-Grade Tests', *Environmental Health Perspectives* 115(8), 1242–1247.
- Miranda, Marie Lynn, Kim, Dohyeong, Reiter, Jerome, Overstreet Galeano, M. Alicia and Maxson, Pamela.** (2009). 'Environmental contributors to the achievement gap', *Neuro-Toxicology* 30(6), 1019–1024.
- Monbiot, George.** (2013). 'Yes, lead poisoning could really be a cause of violent crime', *The Guardian* .

- Needleman, Herbert L., McFarland, Christine, Ness, Roberta B., Fienberg, Stephen E. and Tobin, Michael J.** (2002). 'Bone lead levels in adjudicated delinquents: A case control study', *Neurotoxicology and Teratology* 24(6), 711–717.
- Needleman, Herbert L, Riess, Julie A, Tobin, Michael J, Biesecker, Gretchen E and Greenhouse, Joel B.** (1996). 'Bone Lead Levels and Delinquent Behavior', *JAMA: The Journal of the American Medical Association* 275(5), 363–369.
- Needleman, Herbert L, Schell, Alan, Bellinger, David, Leviton, Alan and Allred, Elizabeth N.** (1990). 'The long-term effects of exposure to low doses of lead in childhood: an 11-year follow-up report', *New England journal of medicine* 322(2), 83–88.
- Nevin, Rick.** (2000). 'How Lead Exposure Relates to Temporal Changes in IQ, Violent Crime, and Unwed Pregnancy', *Environmental Research* 83(1), 1–22.
- Nevin, Rick.** (2007). 'Understanding international crime trends: The legacy of preschool lead exposure', *Environmental Research* 104(3), 315–336.
- Nilsson, J. Peter.** (2009). 'The long-term effects of early childhood lead exposure: Evidence from the phase-out of leaded gasoline', *Institute for Labour Market Policy Evaluation (IFAU) Work. Pap.*
- Olds, David L, Henderson Jr, Charles R, Kitzman, Harriet J, Eckenrode, John J, Cole, Robert E and Tatelbaum, Robert C.** (1999). 'Prenatal and infancy home visitation by nurses: Recent findings', *The future of Children* pp. 44–65.
- Olds, David L, Kitzman, Harriet, Hanks, Carole, Cole, Robert, Anson, Elizabeth, Sidora-Arcoleo, Kimberly, Luckey, Dennis W, Henderson, Charles R, Holmberg, John, Tutt, Robin A et al.** (2007). 'Effects of nurse home visiting on maternal and child functioning: age-9 follow-up of a randomized trial', *Pediatrics* 120(4), e832–e845.
- Poterba, James M.** (1996), Government intervention in the markets for education and health care: how and why?, in 'Individual and social responsibility: Child care, education, medical care, and long-term care in America', University of Chicago Press, pp. 277–308.
- Prüss-Üstün, Annette, Fewtrell, Lorna, Landrigan, Philip J. and Ayuso-Mateos, José Luis.** (2004), Lead Exposure, in **Majid Ezzati, Alan D. Lopez, Anthony Rodgers and Christopher J.L. Murray.**, ed., 'Comparative Quantification of Health Risks', Vol. 2, World Health Organization, pp. 1495–1542.
- Rau, Tomás, Reyes, Loreto and Urzúa, Sergio S.** (2013), The Long-term Effects of Early Lead Exposure: Evidence from a case of Environmental Negligence, Working Paper w18915, National Bureau of Economic Research.

- Reyes, Jessica W.** (2011), Childhood lead and academic performance in Massachusetts, New England Public Policy Center Working Paper 11-3, Federal Reserve Bank of Boston.
- Reyes, Jessica Wolpaw.** (2007). 'Environmental Policy as Social Policy? The Impact of Childhood Lead Exposure on Crime', *The B.E. Journal of Economic Analysis & Policy* 7(1).
- Reyes, Jessica Wolpaw.** (2015). 'Lead Exposure and Behavior: Effects on Antisocial and Risky Behavior among Children and Adolescents', *Economic Inquiry* .
URL: <http://dx.doi.org/10.1111/ecin.12202>
- Ris, M Douglas, Dietrich, Kim N, Succop, Paul A, Berger, Omer G and Bornschein, Robert L.** (2004). 'Early exposure to lead and neuropsychological outcome in adolescence', *Journal of the International Neuropsychological Society* 10(02), 261–270.
- Sandel, Megan, Baeder, Andrea, Bradman, Asa, Hughes, Jack, Mitchell, Clifford, Shaughnessy, Richard, Takaro, Tim K and Jacobs, David E.** (2010). 'Housing interventions and control of health-related chemical agents: a review of the evidence', *Journal of Public Health Management and Practice* 16(5), S24–S33.
- Schnaas, Lourdes, Rothenberg, Stephen J., Flores, Maria-Fernanda, Martinez, Sandra, Hernandez, Carmen, Osorio, Erica, Velasco, Silvia Ruiz and Perroni, Estela.** (2006). 'Reduced Intellectual Development in Children with Prenatal Lead Exposure', *Environmental Health Perspectives* 114(5), 791–797.
- Troesken, Werner.** (2008). 'Lead water pipes and infant mortality at the turn of the twentieth century', *Journal of Human Resources* 43(3), 553–575.
- Wright, John Paul, Dietrich, Kim N, Ris, M. Douglas, Hornung, Richard W, Wessel, Stephanie D, Lanphear, Bruce P, Ho, Mona and Rae, Mary N.** (2008). 'Association of Prenatal and Childhood Blood Lead Concentrations with Criminal Arrests in Early Adulthood', *PLoS Med* 5(5), e101.
- Yeoh, Berlinda, Woolfenden, Susan, Wheeler, Danielle M, Alperstein, Garth and Lanphear, Bruce.** (2009). 'Cochrane review: Household interventions for prevention of domestic lead exposure in children', *Evidence-Based Child Health: A Cochrane Review Journal* 4(2), 951–999.
- Zhang, Nanhua, Baker, Harolyn W, Tufts, Margaret, Raymond, Randall E, Salihu, Hamisu and Elliott, Michael R.** (2013). 'Early childhood lead exposure and academic achievement: evidence from Detroit public schools, 2008–2010', *American Journal of Public Health* 103(3), e72–e77.


9 Figures

Figure 1: Elevated blood lead level intervention policy of the Children's Environmental Health branch within the North Carolina Department of Health

Interpretation of Screening Test Results and Recommended Follow-up	
Blood Lead Level ($\mu\text{g}/\text{dL}$)	Comments
<10	A child with this Blood Lead Level (BLL) is not considered to have an elevated level of exposure. Reassess or rescreen in one year. No additional action is necessary unless exposure sources change.
10-14	The CDC considers 10 $\mu\text{g}/\text{dL}$ to be a level of concern. Perform diagnostic test on venous blood within three months. If the diagnostic test is confirmatory, the child should have follow-up tests at three month intervals until the BLL is <10 $\mu\text{g}/\text{dL}$. Provide family lead education. Refer for nutrition counseling.
15-19	A child in this category should also receive a diagnostic test on venous blood within three months. If the diagnostic test is confirmatory, the child should have additional follow-up tests at three month intervals. Children with this level of exposure should receive clinical management. Parental education and nutritional counseling should be conducted. A detailed environmental history should be taken to identify any obvious sources of lead exposure.
20-44	A child with a BLL in this range should receive a confirmatory venous test within one week to one month. The higher the screening test, the more urgent the need for a diagnostic test. If the diagnostic test is confirmatory, coordination of care and clinical management should be provided. An abdominal x-ray is completed if particulate lead ingestion is suspected. Nutrition and education interventions, a medical evaluation, and frequent retesting (every 3 months) should be conducted. Environmental investigation and lead hazard control is needed for these children.
45-69	A child in this category should receive a confirmatory venous test within 48 hours. If the screening blood lead level is between 60-69 $\mu\text{g}/\text{dL}$, the child should have a venous blood lead level within 24 hours. If confirmatory, case management and clinical management should begin within 48 hours. Environmental investigation and lead hazard control should begin as soon as possible. A child in this exposure category will require chelation therapy and an abdominal x-ray is completed if particulate lead ingestion is suspected.
≥ 70	A child with a BLL ≥ 70 requires immediate hospitalization as lead poisoning at this level is a medical emergency. Confirmatory venous testing should be done as soon as possible. An abdominal x-ray is completed if particulate lead ingestion is suspected and chelation therapy should begin immediately. Case and clinical management including nutrition, education, medical and environmental interventions, must take place as soon as possible.
Information from Centers for Disease Control and Prevention. Screening Young Children for Lead Poisoning: Guidance for State and Local Public Health Offices. November 1997. Atlanta, Georgia. United States Department of Health and Human Services, Public Health Services, CDC, 1997 and Centers for Disease Control and Prevention. Managing Elevated Blood Lead Levels Among Children: Recommendations from the Advisory Committee on Childhood Lead Poisoning Prevention. March 2002	


Notes: This guide represents NC Health Department Policies in 2002 (entirely based on CDC recommendations). Since some of our sample is tested prior to 2002, we have investigated and found no changes in lead policy in the years preceding. Conversations with the NC Childhood Lead Poisoning Prevention Program have confirmed that these guidelines were used at least back to 1991. Based on conversations with health workers in North Carolina and specifically Mecklenburg County, NC, along with inspection of the recommended interventions, the thresholds for which policy is substantially different is the 10 $\mu\text{g}/\text{dL}$ and the 20 $\mu\text{g}/\text{dL}$ threshold. We add emphasis of interventions triggered by underlining the intervention components (excluding further testing).

Figure 2: Observations by First and Second BLL Test




This figure provides a grid with 1st and 2nd BLL test result values indicating treatment and control regions and highlights the variation in BLL between the first and second BLL test as well as the number of observations in our estimation sample for various combinations of first and second BLL test results.

Figure 3: Average Education Outcomes by Blood Lead Level



Notes This figure depicts mean outcomes by the level of initial BLL test result for each of the the education outcomes. The group "NO INTERVENTION, BLL > 10" includes any individual with an initial test over 10µg/dL but without a second test over 10µg/dL. This group includes our control group (1st test >10, 2nd test >5 but <10) as well as other individuals with at least one BLL test >10 but not in our intervention group. Means for our control group are, in general, worse off compared to this group and are listed in Table 2. The "INTERVENTION, BLL > 10" group represents the treatment group in our estimation sample—those who are eligible for the intervention following two consecutive tests over the threshold of 10µg/dL.


Figure 4: Average Behavioral Outcomes by Blood Lead Level




Notes This figure depicts mean outcomes by the level of initial BLL test result for each of the behavioral outcomes. The group "NO INTERVENTION, BLL > 10" includes any individual with an initial test over 10 μ g/dL but without a second test over 10 μ g/dL. This group includes our control group (1st test >10, 2nd test >5 but <10) as well as other individuals with at least one BLL test >10 but not in our intervention group. Means for our control group are, in general, worse off compared to this group and are listed in Table 2. The "INTERVENTION, BLL > 10" group represents the treatment group in our estimation sample—those who are eligible for the intervention following two consecutive tests over the threshold of 10 μ g/dL.

Figure 5: Change in BLLs

A. Final BLL for Intervention Group




B. Change in BLL for Intervention Group




Notes: This figure plots the distribution of observations by the final BLL value (A) and the net change in detected BLL between the 2nd and 3rd test (B) for the intervention group (first and second test $\geq 10\mu\text{g/dL}$).

Figure 6: Estimated Treatment Effect Relative to Alternative Control Groups (by Average BLL)

A. Sample = At least one BLL test



B. Sample = At least two BLL tests



This figure plots estimates of eligibility for the intervention on the summary index outcomes comparing our treatment group (two tests with BLL 10 or more) with groups defined by their average BLL test results. These control groups are no longer restricted to having an initial BLL test above 10. Panel A includes individuals with at least one BLL test in the control group while Panel B includes only those with at least two BLL tests. The sample sizes for each alternative control group indicated by the labels on the horizontal axis are as follows: *Panel A*: Avg BLL 1-3 (5,540); Avg BLL 3-5 (7,959); Avg BLL 5-7 (3,987); Avg BLL 7-10 (2,044); Avg BLL 10-15 (638); Avg BLL >15 (145). *Panel B*: Avg BLL 1-3 (783); Avg BLL 3-5 (1,450); Avg BLL 5-7 (681); Avg BLL 7-10 (351); Avg BLL 10-15 (72); Avg BLL >15 (15). All regressions include the full set of control variables listed in table notes for Table 4.

10 Tables

Table 1: Means of demographic, housing, and neighborhood characteristics

	<u>Intervention</u>	<u>Control</u>	<u>Difference</u>
<u>Background Characteristics</u>			
Male	0.61 (0.49)	0.58 (0.49)	0.02 (0.06)
Minority	0.77 (0.42)	0.77 (0.42)	-0.00 (0.05)
Stand Alone Residence	0.58 (0.50)	0.57 (0.50)	-0.00 (0.06)
Home Built pre 1978	0.79 (0.41)	0.78 (0.42)	-0.01 (0.05)
Past Lead Tests at a Home (mean $\mu\text{g}/\text{dL}$)	4.40 (1.16)	4.52 (1.51)	-0.12 (0.25)
Age at Blood Lead Test (months)	28.07 (17.21)	25.57 (14.15)	2.50 (1.82)
Mother Education (years)	11.92 (2.96)	11.45 (2.28)	0.47 (0.37)
Birth Weight (ozs)	115.09 (20.37)	110.90 (21.57)	4.19 (3.05)
Index of Neighborhood Attributes	-0.48 (0.88)	-0.58 (0.89)	0.10 (0.10)
F-stat (p-value)			0.237
Observations	119	182	301

Notes: This table reports means and standard deviations for the group eligible for intervention (two tests $\geq 10\mu\text{g}/\text{dL}$) and our control group (first test $\geq 10\mu\text{g}/\text{dL}$, second test ≥ 5 but $< 10\mu\text{g}/\text{dL}$) as well as the mean difference and the standard error of the difference. Any statistically significant differences are noted with * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. The number of observations by Intervention and Control groups for each variable is reported at the bottom with the exception of the following variables: *Stand Alone Residence* (99/143); *Built Pre 1978* (108/148); *Past Lead Tests at a Home* (49/87); *Mother Education* (76/131); *Birthweight* (76/131). The *Index of Neighborhood Attributes* is an index measure of disadvantage derived from census block group variables (based on address at first lead test). We calculate this summary measure from an unweighted z-score sum of the percent of households in a 2000 Census Block Group (CBG) without a high school graduate, the CBG poverty rate, the CBG fraction of single female headed households, and the CBG population density. The p-value from a F-test of joint-significance of all of the background characteristics is also reported (see Table A4 for the balance test results).

Table 2: Means of education and behavior outcomes

	<u>Intervention</u>	<u>Control</u>	<u>Difference</u>
Blood lead level at initial test ($\mu\text{g}/\text{dL}$)	17.85 (8.25)	12.09 (4.41)	5.76*** (0.73)
Education Index	0.08 (0.60)	-0.05 (0.71)	0.13* (0.08)
Reading Test Score (avg 3-5th grade)	-0.44 (0.83)	-0.58 (0.91)	0.14 (0.12)
Math Test Score (avg 3-5th grade)	-0.46 (0.81)	-0.53 (0.96)	0.07 (0.12)
Repeat a grade (grades 1-5)	0.15 (0.36)	0.14 (0.35)	0.01 (0.04)
Reading Test Score (avg 6-8th grade)	-0.32 (0.81)	-0.50 (0.95)	0.18 (0.12)
Math Test Score (avg 6-8th grade)	-0.31 (0.82)	-0.43 (0.88)	0.12 (0.11)
Repeat a grade (grades 6-9)	0.14 (0.35)	0.21 (0.41)	-0.07 (0.05)
Adolescent Antisocial Behavior Index	-0.15 (0.47)	0.10 (0.83)	-0.25*** (0.08)
Days Suspended (6th-10th grade)	9.25 (15.80)	17.67 (32.44)	-8.42*** (3.20)
Days Absent (6th-10th grade)	30.61 (36.31)	45.65 (54.71)	-15.05*** (5.70)
School Reported Crimes (6th-10th grade)	1.97 (3.40)	3.45 (6.75)	-1.47** (0.67)
Ever Arrested	0.08 (0.27)	0.18 (0.38)	-0.10** (0.04)
Ever Arrested - Violent	0.03 (0.16)	0.12 (0.32)	-0.09*** (0.03)
Ever Arrested - Property	0.04 (0.20)	0.07 (0.26)	-0.03 (0.03)
Observations	119	182	301

Notes: This table reports means and standard deviations of blood lead levels and outcome variables for the group eligible for intervention (two tests $\geq 10\mu\text{g}/\text{dL}$) and our control group (first test $\geq 10\mu\text{g}/\text{dL}$, second test ≥ 5 but $< 10\mu\text{g}/\text{dL}$) as well as the mean difference between the two groups along with the standard error of the difference. Any statistically significant differences in the means are noted with * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. We follow the methodology in [Anderson \(2008\)](#) to create a summary index (a weighted mean of standardized outcomes). The education index includes 3rd through 5th grade math and reading test score results and grade retention between 3rd and 9th grade. All test scores are standardized based on state-wide averages by grade and calendar year. The antisocial behavior index includes measures of number of days suspended and absences (6th through 10th grade), school reported crimes, and criminal arrests between the ages of 16 and 18.

Table 3: Do individuals comply with the elevated BLL intervention?

	(1) Had 3rd BLL Test w/in 100 days	(2) Had 3rd BLL Test	(3) Total # of BLL Tests	(4) Months b/t 2nd & 3rd Test	(5) Future Lead Remed- iation
Intervention	0.438*** (0.045)	0.524*** (0.045)	2.661*** (0.318)	-7.563*** (1.363)	0.048 (0.032)
Dep. Var. (mean) for Control Group	0.08	0.23	2.35	12.22	0.03
Observations	301	301	301	113	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This table presents results for specifications with dependent variables assessing whether individuals eligible for an intervention appear to comply with the recommended procedures including (1) whether they show up for follow up test following the second confirmatory test within 100 days (2) whether they ever show up for a third test, (3) the total number of BLL tests, (4) the timing between the follow up tests, and (5) whether the property is remediated following a referral to the LeadSafe Charlotte remediation program. All regressions include the full set of control variables listed in the table notes of Table 4. There are fewer observations for column (4) due to the limited number of individuals that have third tests.

Table 4: Effects of an elevated BLL intervention on education and behavioral outcomes

	(1) Education Index	(2) Reading (3-5th)	(3) Math (3-5th)	(4) Repeat Grade (1-5th)	(5) Reading (6-8th)	(6) Math (6-8th)	(7) Repeat Grade (6-9th)
Intervention	0.117* (0.067)	0.153 (0.119)	0.099 (0.117)	0.035 (0.039)	0.219** (0.102)	0.163 (0.102)	-0.036 (0.043)
Observations	301	240	244	301	235	236	301

	(1) Adolescent Antisocial Behavior Index	(2) Days Suspended (6-10th)	(3) Days Absent (6-10th)	(4) School Crimes (6-10th)	(5) Ever Arrested	(6) Ever Arrested Violent	(7) Ever Arrested Property
Intervention	-0.184** (0.082)	-5.936** (2.698)	-9.786** (4.281)	-1.219** (0.607)	-0.073* (0.043)	-0.076*** (0.027)	-0.017 (0.037)
Observations	301	301	301	301	301	301	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This table reports the estimated effects of EBLL intervention eligibility on the various educational and behavioral outcomes. The treatment and control groups as well as the outcome variables are defined in Table 2. All estimates in this table are based off of a regression specification including the following controls: indicators for gender, minority (black or hispanic), birth year, single family home, pre-1978 parcel age, missing school test scores, missing birth record, missing parcel information; and continuous controls for birthweight, age at blood test, average previous lead levels for prior households in the home, mother's years of education, and a CBG-based neighborhood index described in Table 1.

Table 5: Heterogeneous Effects by Intensity of Intervention

	(1)	(2)
	Education Index	Adolescent Antisocial Behavior Index
Intervention (20+)	0.295* (0.161)	-0.276** (0.121)
Intervention (15+)	-0.069 (0.124)	0.056 (0.110)
Intervention (10+)	0.068 (0.073)	-0.152 (0.094)
Observations	301	301
Intervention*Large Drop in BLL	0.102 (0.135)	-0.223** (0.095)
Intervention	0.091 (0.069)	-0.128 (0.086)
Observations	301	301
Intervention*Quick Time to Next BLL Test	0.107 (0.172)	-0.080 (0.127)
Intervention	0.105 (0.069)	-0.175** (0.082)
Observations	301	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

This table presents results by different measures of the intensity of the potential intervention. All regressions include the full set of control variables listed in the table notes of Table 4. In the top panel of this table, we include indicators for potentially higher intensity treatment categories based on thresholds outlined in CDC recommendations summarized in Figure 1. We create indicators for those within the treatment group who have a test above $15\mu\text{g/dL}$ and those with a test above $20\mu\text{g/dL}$. Note that these indicators are not mutually exclusive; An individual with a confirmatory test over 20 would have each of the three treatment level indicators equal to one. For the second panel and third panel of the table, we test for heterogeneous effects for other measures potentially capturing the intensity of the response to confirmed elevated blood lead levels. In the second panel, we define large drop as those individuals that see a drop in BLL of more than 5 BLL between 2nd and 3rd test. In the third panel, we define quick time between 2nd and 3rd test based on less than 1 month between 2nd (confirmatory) test and a 3rd BLL test.

Table 6: Educational and Behavioral Differences for Prior and Future Residents

	(1)	(2)
	Education Index	Adolescent Antisocial Behavior Index
<u>Prior Residents</u>		
Intervention Parcel	0.030 (0.049)	0.001 (0.047)
Observations	1,363	1,363
<u>Future Residents</u>		
Intervention Parcel	0.100 (0.076)	-0.133 (0.093)
Observations	430	430

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This tables reports the estimated difference in the summary index outcomes for children not included in our analysis but who were living in the treatment and control parcels either before or after the children included in our estimation sample. The first panel presents results for individuals that lived at the same address *prior* to our sample of treatment and control observations; while the second panel presents results for individuals living at the same address *after* our estimation sample. We drop any parcels that contain both treatment and control observations. All regressions include the full set of control variables listed in the table notes of Table 4.

Table 7: Estimated Costs of Elevated BLL Intervention

	Cost	Pr(Cost Incurred)	Estimated Cost
<u>Medical Costs</u>			
Doctor's Visit/Nutritional Assessment	\$250	1	\$250
Additional BLL Tests ($\times 3$)	\$225	1	\$225
<u>Home Inspection and Remediation</u>			
Home inspection	\$650	1	\$650
Lead-based paint remediation	\$7,300	0.073	\$533
<u>Social Services Cost</u>			
Child Development Services/Case Management (3yrs)	\$9,000	0.403	\$3,630
Total estimated cost			\$5,288

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Cost figures and most probability calculation based on detailed conversations with social service providers in Mecklenburg County, NC. Probabilities and costs of lead testing come from the NC Children's Environmental Health Branch of the Department of Health and Human Services. Information on the costs of social services is from Mecklenburg County Children's Developmental Services. Information on lead remediation costs is based on HUD grant reporting records from Leadsafe Charlotte. Leadsafe remediation probability based on estimated probabilities for our intervention group given by Table 3.

Appendix

Background on Data Sources and Sample Construction

Our primary source of data is the blood lead surveillance data from the state registry maintained by the NC Childhood Lead Poisoning Prevention Program of the Children's Environmental Health Branch. This dataset includes a child's name, gender, birth date, test date, blood lead level (BLL) and home address. The North Carolina State Laboratory for Public Health (Raleigh, NC) conducted 90 percent of the lead analyses of the blood samples and all BLL values are stored as integers with a value of $1\mu\text{g}/\text{dL}$ (micrograms per deciliter) given to children without any detectable lead.

Our analysis focuses only on children living in Mecklenburg County and includes all BLL tests for a child between 1993 and 2008. North Carolina requires all children participating in Medicaid or the Special Nutrition Program for Women, Infants and Children (WIC) to be screened for lead at 1 or 2 years of age. Other children are screened if a parent responds "yes" or "don't know" to any of the questions on a CDC Lead Risk Assessment Questionnaire. The North Carolina Blood Lead Surveillance Group estimates that it screened between 21.9 and 30.4% of children one and two years of age from 1995 through 1998 and we expect screening rates were similar during our analysis period (Miranda et al. (2007)). This dataset provides multiple blood lead level tests per child which allows us to determine which children received various lead policy interventions due to two tests with BLL of $10\mu\text{g}/\text{dL}$ or above.

We subsequently match individual children to two additional databases in order to examine the impact of interventions on educational and behavioral outcomes. All matches are conducted using first and last name as well as date of birth and we incorporate fuzzy matches for names in some cases. Our first database is the administrative records from Charlotte-Mecklenburg Schools (CMS) that span kindergarten through 12th grade and the school years 1998-1999 through 2010-2011. This dataset includes each student that attended a public school in the City of Charlotte for at least one semester and provides annual data for each year of matriculation. Specifically, we incorporate student demographics on race and home address, yearly end-of-grade (EOG) test scores for grades 3 through 8 in math and reading, number of days absent, days suspended from school as well as the number of incidents of school crime.⁵⁶ We are able to match 65 percent of lead tests to a student record in CMS. This match rate improves to 74 percent for our policy sample of individuals with two tests and one test $>10\mu\text{g}/\text{dL}$.

In order to examine adult criminal outcomes we match our lead database to a registry of all adult (defined in North Carolina as age 16 and above) arrests in Mecklenburg County from

⁵⁶According to NC State Statute 115C – 288(g), any incident at school involving any violent or threats of violent behavior, property damage, theft or drug possession must officially be reported to the NC school crimes division. This statute ensures that this measure of school crime is consistently reported across schools and cannot be treated differently based on school administrators.

2006 to 2013. We use first name, last name and date of birth to link individuals across the two data sources. While over 90 percent of the matches are exact, we recover additional matches using an algorithm for partial matches that has been used and validated in Deming (2011). The Mecklenburg County Sheriff (MCS) tracks arrests and incarcerations across individuals using a unique identifier that is established with fingerprinting. The arrest data include information on the number and nature of charges as well as the date of arrest. This data allows us to observe adult criminality regardless of whether a child later transferred or dropped out of CMS schools with the main limitation being that it only includes crimes committed within Mecklenburg County. The quality of matching between the lead and arrests databases is not directly measurable since one cannot distinguish between those lead tested individuals never arrested versus individuals who do not match due to clerical errors in names or moving out of the county. We can speak to the quality of matches using the arrest database by the fact that we are able to match approximately 94 percent of arrest records for a given cohort to our CMS education database.

In order to provide some basic controls for parental and housing factors, we draw on two additional databases. The first database is the universe of birth certificate records from the state of North Carolina from 1990-2002. As with previous databases, we are able to match our lead database to the birth records database using name and date of birth. In the case of birth records we are primarily interested in two variables, birth weight and mother's years of education.⁵⁷ We are able to match approximately 54 percent of birth records to our lead database. Even though this match rate is somewhat lower than our other databases, the variables from this database are simply used as control variables and we later show that this match rate is unrelated to our analysis of lead policy interventions. The second database is county assessor's data for all parcels on an annual basis from 2002-2012 in Mecklenburg County, NC. For this database, we match our lead data to parcel records based on home address given for an individual's first lead test. We augment this parcel data with building permits for all home renovations from 1995-2012. This database on parcels allows us to generate variables for prior home renovations, age and type of housing structure. The lead database is matched to parcels records 86 percent of the time with differences primarily a result of incomplete homes address information.

In some of our analysis, we merge into our dataset two additional data elements. First, we merge data from the LeadSafe Charlotte program which contains detailed data on the addresses of approximately 2,500 homes (single-family and multi-family) which have been lead inspected or lead remediated and certified lead safe since 1998. We match LeadSafe addresses to our county parcel data based on parcel addresses with 20 LeadSafe homes unable to be successfully matched to parcel records. Second, we construct a measure of siblings using birth records data. In order to be characterized as a sibling, two individuals must share a mother's first name, last

⁵⁷We focus on maternal education because paternal information is not consistently reported on birth records in NC during this time period.

name and date of birth based on Mecklenburg County birth records.

Summary Index Construction

We follow the methodology in [Anderson \(2008\)](#) to create two summary index outcome measures: educational performance and adolescent antisocial behavior. The antisocial behavior index is created to include measures of number of days suspended and unapproved absences (6th through 10th grade), school reported crimes, and criminal arrests between the ages of 16 and 18. The education index includes 3rd through 8th grade math and reading test score results and grade retention between 1st and 9th grade.

A summary of the steps to create an index are listed below. See [Anderson \(2008\)](#) for additional detail in calculation of a summary index.

1. Switch signs where necessary so the positive direction indicates a larger outcome effect.
2. Demean outcomes and convert to effect sizes by dividing by its control group standard deviation.
3. Define groupings of outcomes.
4. Create a new variable that is a weighted average of the outcomes in each grouping. When constructing the weighted average, weight each element by the inverse of the covariance matrix of the standardized outcomes in each group.
5. Regress the new weighted average for each group on intervention status to estimate treatment effects.

Table A1: Variable Definitions

Variable Name	Description	Source
Outcomes		
<i>Education Index</i>	An index with mean zero and standard deviation one based on the academic outcomes listed below	CMS pupil records 1999-2011
Reading Test Score	The average reading test scores for given grades All test scores are normalized to mean zero and standard deviation one based on all NC tests in the same grade and testing year	
Math Test Score	The average math test scores for given grades	
Repeat a Grade	Indicator if a student ever repeats a given school grade.	
<i>Antisocial Behavior Index</i>	An index with mean zero and standard deviation one based on the behavioral outcomes listed below	
Days Suspended	Total days suspended (in + out of school) (6th-10th grade)	CMS pupil records 1999-2011
Days Absent	Total days absent (6th-10th grade)	CMS pupil records 1999-2011
School Crimes	Total crimes reported for a given student (6th-10th grade)	CMS pupil records 1999-2011
Ever Arrested (16-18 years old)	Indicator if ever arrested for a crime in Mecklenburg County/Charlotte NC	Mecklenburg County Sheriff 1998-2014
Ever Arrested - Violent Crime	Indicator if Ever Arrested for a violent crime (assault, manslaughter, robbery)	Mecklenburg County Sheriff 1998-2014
Ever Arrested - Property Crime	Indicator if Ever Arrested for a property crime (burglary, larceny, forgery/fraud, auto theft)	Mecklenburg County Sheriff 1998-2014
Background Characteristics		
Male	Indicator for gender	CMS pupil records 1999-2011
Minority	Indicator for race equal to black or hispanic	CMS pupil records 1999-2011
Stand Alone Residence*	Indicator for single unit housing structure (single-family residence)	Mecklenburg County Assessor 1999-2015
Home Built pre-1978*	Indicator for home built before 1978	Mecklenburg County Assessor 1999-2015
Past Lead Tests at a Home (mean BLL)*	Average of all prior BLL tests with the same residential address as a given observation	NC Blood Surveillance 1993-2008
Age at BLL Test	Age in months for initial BLL test	NC Blood Surveillance 1993-2008
Mother's Education*	Years of education reported for a child's mother	NC Birth Records 1990-2008
Birth Weight*	An observation's birth weight (ozs)	NC Birth Records 1990-2008
Index of Neighborhood Attributes	Average of standardized values (mean zero, standard deviation one) for four Census Block Group neighborhood variables: Percent in Poverty; Percent of adults with no high school diploma; Percent of households with single female-headed households; Population density	US Census 2000

* Some values of these variables are missing for our main estimation sample. For estimation, any observations with missing values are assigned a zero value and we create an indicator for missing value for that variable.

Table A2: Means of demographic, housing, and neighborhood characteristics

	All Students	Lead Tested	BLL 5-9	BLL ≥10
<u>Background Characteristics</u>				
Male	0.51 (0.50)	0.51 (0.50)	0.52 (0.50)	0.55 (0.50)
Minority	0.49 (0.50)	0.60 (0.49)	0.69 (0.46)	0.70 (0.46)
Stand Alone Residence	0.67 (0.47)	0.65 (0.48)	0.63 (0.48)	0.66 (0.48)
Home Built pre 1978	0.43 (0.49)	0.65 (0.48)	0.72 (0.45)	0.74 (0.44)
Past Lead Tests at a Home (mean $\mu\text{g/dL}$)	3.91 (1.21)	4.09 (1.16)	4.20 (1.18)	4.43 (1.52)
Age at Blood Lead Test	2.12 (1.50)	2.20 (1.53)	2.15 (1.42)	1.89 (1.26)
Mother Education (years)	13.28 (2.48)	12.69 (2.52)	12.33 (2.44)	12.08 (2.40)
Birth Weight (ozs)	115.81 (21.86)	113.52 (21.95)	112.54 (21.39)	111.22 (20.56)
Index of Neighborhood Attributes	0.08 (0.76)	-0.28 (0.85)	-0.42 (0.86)	-0.44 (0.87)
Observations	153,039	19,731	5,857	935

This table reports means and standard deviations for variables given in Table 1 for the full population of public students as well as by different lead testing BLL values for individuals born between 1990 and 1997.

Table A3: Means of education and behavior outcomes

	All Students	Lead Tested	BLL 5-9	BLL ≥10
Blood lead level (µg/dL)	4.144 (3.115)	4.220 (3.236)	6.169 (1.245)	13.129 (7.900)
<u>Education Outcomes</u>				
Reading Test Score (avg 3-5th grade)	-0.030 (0.965)	-0.204 (0.956)	-0.364 (0.934)	-0.474 (0.916)
Math Test Score (avg 3-5th grade)	-0.033 (0.973)	-0.205 (0.953)	-0.366 (0.921)	-0.427 (0.918)
Repeat a grade (grades 1-5)	0.046 (0.210)	0.102 (0.303)	0.133 (0.339)	0.140 (0.347)
Reading Test Score (avg 6-8th grade)	-0.033 (0.967)	-0.174 (0.952)	-0.335 (0.932)	-0.409 (0.920)
Math Test Score (avg 6-8th grade)	-0.038 (0.969)	-0.175 (0.935)	-0.324 (0.888)	-0.378 (0.888)
Repeat a grade (grades 6-9)	0.101 (0.302)	0.142 (0.349)	0.193 (0.395)	0.197 (0.398)
<u>Antisocial Behavior Outcomes</u>				
Days Suspended (6-10th grade)	4.34 (13.39)	8.49 (19.85)	11.29 (22.88)	14.35 (26.75)
Days Absent (6-10th grade)	20.78 (31.00)	30.64 (39.30)	37.23 (45.74)	41.31 (47.65)
School Reported Crimes (6-10th grade)	0.93 (3.02)	1.96 (4.63)	2.44 (5.09)	2.77 (5.40)
Ever Arrested	0.05 (0.21)	0.08 (0.27)	0.11 (0.31)	0.12 (0.33)
Ever Arrested - Violent	0.02 (0.13)	0.04 (0.18)	0.05 (0.21)	0.06 (0.24)
Ever Arrested - Property	0.02 (0.14)	0.04 (0.19)	0.05 (0.22)	0.06 (0.24)
Observations	153,039	19,731	5,857	935

This table reports means and standard deviations for variables given in Table 2 for the full population of public students as well as by different lead testing BLL values for individuals born between 1990 and 1997. *Note:* The mean blood lead level for All Students does not equal the mean blood lead level for the Lead Tested individuals since some students are not matchable to lead testing data.

Table A4: Balancing test for observable characteristics

	(1) Intervention (2 tests 10+)
Male	-0.008 (0.058)
Minority	0.083 (0.076)
Home Built pre 1978	0.144** (0.068)
Past Lead Tests at a Home (mean $\mu\text{g}/\text{dL}$)	-0.028 (0.031)
Stand Alone Residence	-0.029 (0.064)
Age at Blood Lead Test (months)	0.002 (0.002)
Birth Weight (ozs)	0.001 (0.002)
Mother Education (years)	0.023* (0.013)
Index of Neighborhood Attributes	0.047 (0.037)
F-Stat (p-value)	0.237
Observations	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

This table presents results from a balance test in which the dependent variable is an indicator equal to one if individual received two tests $\geq 10\mu\text{g}/\text{dL}$. The reported p-value at the bottom of each panel is the result of an F-test of joint-significance of all of the reported variables above. The regression also includes year-of-birth indicators and indicator variables for no match to a birth record or a parcel record.

Table A5: Does the elevated BLL intervention affect matching to data sets or residential mobility?

	<u>Intervention</u>	<u>Control</u>	<u>Difference</u>
Matched to CMS record (incl. in est. sample)	0.76 (0.43)	0.78 (0.41)	-0.022 (0.043)
Observations	156	232	388
Did Not Enroll in CMS Secondary School	0.11 (0.32)	0.09 (0.29)	0.019 (0.037)
Observations	108	174	280
Parcel Info Missing	0.17 (0.38)	0.21 (0.41)	-0.046 (0.047)
Birth Record Missing	0.36 (0.48)	0.28 (0.45)	0.081 (0.055)
Change in Residence btw Test and School	0.64 (0.48)	0.72 (0.45)	-0.086 (0.058)
Observations	119	182	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

This table reports means and standard deviations of data matching and mobility indicators for the group eligible for intervention (two tests $\geq 10\mu\text{g/dL}$) and our control group (first test $\geq 10\mu\text{g/dL}$, second test ≥ 5 but $< 10\mu\text{g/dL}$) as well as the mean difference and the standard error of the difference. Any statistically significant differences are noted with * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. The first row presents a comparison across the two groups of the probability of an individual meeting the BLL testing criteria being matched to a CMS school record (which is required for inclusion in our estimation sample). The second row compares rates (conditional on primary school enrollment) of not enrolling in a CMS secondary school to test for differential attrition rates for adolescent outcomes. The third and fourth rows compare rates of matching records to the birth record and parcel characteristic databases for those in our estimation sample. Finally, the fifth row compares rates of residential mobility between the first blood lead test and CMS school enrollment for our estimation sample. An indicator for a change in residence is created through a comparison of the residential address listed in each database.

Table A6: Effects of eligibility for the elevated BLL intervention adding sets of controls

	(1) Education Index	(2) Adolescent Antisocial Behavior Index
<u>1. Uncontrolled mean differences</u>		
Intervention	0.129* (0.068)	-0.247*** (0.081)
<u>2. Add controls for bith year</u>		
Intervention	0.142** (0.070)	-0.244*** (0.080)
<u>3. Add controls for individual characteristics</u>		
Intervention	0.138** (0.066)	-0.221*** (0.075)
<u>4. Add controls for age at BLL test</u>		
Intervention	0.136** (0.066)	-0.219*** (0.075)
<u>5. Add controls for maternal educ</u>		
Intervention	0.116* (0.066)	-0.199** (0.077)
<u>6. Add controls for parcel characteristics</u>		
Intervention	0.125* (0.068)	-0.194** (0.081)
<u>7. Add controls for nbhd characteristics</u>		
Intervention	0.117* (0.067)	-0.184** (0.082)
P-value (model 1 = model 7)	0.47	0.11
Observations	301	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This table reports the estimated coefficient on an indicator for eligibility for the BLL intervention for specifications which incrementally add sets of control variables as indicated by the row title. All regressions are based on the primary estimation sample with the control and intervention groups defined as in Table 4. The first row reports the mean uncontrolled differences as previously reported in Table 2. The following controls are added in subsequent rows: (2) indicators for birth year and an indicator for whether there are any missing outcomes in the creation of the index variable; (3) birthweight and indicators for missing birth record, gender, and minority (black or Hispanic) status; (4) age at the time of the initial BLL test; (5) years of maternal education ; (6.) single family home indicator, pre-1978 indicator, building age, blood lead lead levels for previous children tested in home ; (7) Index of Neighborhood Attributes (an unweighted z-score sum of the percent of households without a high school graduate, the CBG poverty rate, the fraction of single female headed households, and the CBG population density).

Table A7: Effects of an elevated BLL intervention on summary index outcomes: *Robustness Check - Different Control Variables*

	(1) Education Index	(2) Adolescent Antisocial Behavior Index
<u>1. Main Results</u>		
Intervention	0.117* (0.067)	-0.184** (0.082)
<u>2. Add control for test type</u>		
Intervention	0.115 (0.070)	-0.185** (0.079)
<u>3. Control for initial BLL fixed effects</u>		
Intervention	0.055 (0.077)	-0.150 (0.110)
<u>4. Control for days between tests</u>		
Intervention	0.096 (0.078)	-0.170** (0.081)
Observations	301	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This table presents results from specifications which add various controls to test for robustness. All regressions include the full set of control variables listed in the table notes of Table 4 and the primary estimation sample with the control and intervention groups defined as in Table 4. The first panel reports our primary estimation results for comparison purposes. The second panel includes an indicator for the type of BLL test used for a second (confirmatory) test. The majority of tests are capillary specimens typically obtained by the a finger prick while some follow up tests are a venous blood draw. The third panel of this table presents results from a specification which includes indicator variables for each initial BLL value to assess whether there may be substantial selection concerns arising from parents responding differently to initial results. Finally, the fourth panel presents results controlling for the days between the initial and follow-up confirmatory test. This variable is correlated with treatment assignment through a mechanical mechanism: due to the 30 day half-life of lead in the blood, a quicker follow-up is more likely to yield a higher confirmatory test result.

Table A8: Regression Discontinuity Results

	(1)	(2)
	Education Index	Adolescent Antisocial Behavior Index
<u>1. Full Sample RDD, BW=5-14</u>		
Intervention	0.071 (0.090)	-0.228** (0.099)
Observations	6,575	6,575
<u>2. Estimation Sample RDD, 2nd BLL Test BW=5-14</u>		
Intervention	-0.163 (0.153)	-0.163 (0.188)
Observations	248	248
<u>3. Estimation Sample RDD, any 2nd BLL Test</u>		
Intervention	-0.062 (0.200)	-0.228 (0.336)
Observations	301	301

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

The top panel presents results from a local linear regression discontinuity design (RDD) using the initial BLL test result as a running variable and thus utilizing all students with an initial BLL from 5 μ g/dL through 15 μ g/dL. The second and third panels present results from a local linear RDD using the second (confirmatory) BLL test result as the running variable. The second panel restricts the bandwidth again to those with a 2nd test result of 5 through 14 μ g/dL. The third panel allows data from the entire estimation sample and estimates an RDD again using the confirmatory test as the running variable but without any restriction on the bandwidth. All regressions include the full set of control variables listed in the table notes of Table 4.


Table A9: Effects of an elevated BLL intervention on summary index outcomes for siblings

	(1)	(2)
	Education Index	Adolescent Antisocial Behavior Index
<u>Younger Siblings</u>		
Younger Sibling of Child (>10 , >10)	0.091 (0.170)	-0.216 (0.241)
Observations	120	120
<u>Older Siblings</u>		
Older Sibling of Child (>10 , >10)	-0.009 (0.278)	-0.123 (0.262)
Observations	88	88

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Standard errors robust to arbitrary within-CBG correlation in parentheses.

Notes: This table presents results for a sample of siblings of our estimation sample. Due to small sample sizes, we compare siblings between the intervention group and the broader control group defined by individuals whose first BLL test result was $\geq 10\mu\text{g/dL}$. We limit our analysis to siblings within 3 years of age. Siblings are defined based on being born to the same mother (identified by first name, last name and date of birth). All regressions include the full set of control variables listed in the table notes of Table 4.


Figure A7: Outcomes by BLL




Notes: This figure provides plots average index outcomes by BLL values in 2 μ g/dL bins. The top panel (A1 and B1) plots outcomes for all children with BLL test results based on the value of their first test result. The second panel (A2 and B2) includes only those in our estimation sample and plots outcomes by the second (confirmatory) BLL test result horizontal variable.

Figure A8: Blood Lead Testing Variation

A: Distribution of all BLL test results



B: Distribution of BLL test results - only children with at least 2 BLL tests



Notes: Panel A of the figure provides the distribution of all first BLL tests in comparison to those individuals that ever had a second BLL test for the full blood surveillance dataset. Panel B further restricts this comparison such that both distributions only contain individuals with two BLL tests.